

## JEST 2017

### Part-A: 1-Mark Questions

Q1. A thin air film of thickness  $d$  is formed in a glass medium. For normal incidence, the condition for constructive interference in the reflected beam is (in terms of wavelength  $\lambda$  and integer  $m = 0, 1, 2, \dots$ )


(a)  $2d = (m - 1/2)\lambda$

(b)  $2d = m\lambda$

(c)  $2d = (m - 1)\lambda$

(d)  $2\lambda = (m - 1/2)d$

Q2. Consider the circuit shown in the figure where  $R_1 = 2.07 k\Omega$  and  $R_2 = 1.93 k\Omega$ . Current source  $I$  delivers  $10mA$  current. The potential across the diode  $D$  is  $0.7V$ . What is the potential at  $A$ ?



(a)  $10.35V$

(b)  $9.65V$

(c)  $19.30V$

(d)  $4.83V$

Q3.  $\int_{-\infty}^{+\infty} (x^2 + 1) \delta(x^2 - 3x + 2) dx = ?$

(a) 1

(b) 2

(c) 5

(d) 7

Q4. A bead of mass  $M$  slides along a parabolic wire described by  $z = 2(x^2 + y^2)$ . The wire rotates with angular velocity  $\Omega$  about the  $z$ -axis. At what value of  $\Omega$  does the bead maintain a constant nonzero height under the action of gravity along  $-\hat{z}$ ?

(a)  $\sqrt{3g}$

(b)  $\sqrt{g}$

(c)  $\sqrt{2g}$

(d)  $\sqrt{4g}$

Q5. Which one is the image of the complex domain  $\{z | xy \geq 1, x + y > 0\}$  under the mapping  $f(z) = z^2$ , if  $z = x + iy$ ?

(a)  $\{z | xy \geq 1, x + y > 0\}$

(b)  $\{z | x \geq 2, x + y > 0\}$


(c)  $\{z | y \geq 2 \forall x\}$

(d)  $\{z | y \geq 1 \forall x\}$

Q6. After the detonation of an atom bomb, the spherical ball of gas was found to be of 15 meter radius at a temperature of  $3 \times 10^5 \text{ K}$ . Given the adiabatic expansion coefficient  $\gamma = 5/3$ , what will be the radius of the ball when its temperature reduces to  $3 \times 10^3 \text{ K}$ ?

- (a) 156m                      (b) 50m                      (c) 150m                      (d) 100m

Q7. What is  $Y$  for the circuit shown below?



- (a)  $Y = \overline{(A + \bar{B})}(\bar{B} + C)$                       (b)  $Y = \overline{(A + \bar{B})}(\bar{B} + C)$   
 (c)  $Y = \overline{(\bar{A} + B)}(\bar{B} + C)$                       (d)  $Y = \overline{(A + B)}(\bar{B} + C)$

Q8. What is the dimension of  $\frac{\hbar \partial \psi}{i \partial x}$ , where  $\psi$  is a wavefunction in two dimensions?

- (a)  $\text{kg m}^{-1} \text{s}^{-2}$                       (b)  $\text{kg s}^{-2}$                       (c)  $\text{kg m}^2 \text{s}^{-2}$                       (d)  $\text{kg s}^{-1}$


Q9. A plane electromagnetic wave propagating in air with  $\vec{E} = (8\hat{i} + 6\hat{j} + 5\hat{k})e^{i(\omega t + 3x - 4y)}$  is incident on a perfectly conducting slab positioned at  $x = 0$ .  $\vec{E}$  field of the reflected wave is


- (a)  $(-8\hat{i} - 6\hat{j} - 5\hat{k})e^{i(\omega t + 3x + 4y)}$                       (b)  $(-8\hat{i} + 6\hat{j} - 5\hat{k})e^{i(\omega t + 3x + 4y)}$   
 (c)  $(-8\hat{i} + 6\hat{j} - 5\hat{k})e^{i(\omega t - 3x - 4y)}$                       (d)  $(-8\hat{i} - 6\hat{j} - 5\hat{k})e^{i(\omega t - 3x - 4y)}$

Q10. Let  $\Lambda = \begin{pmatrix} 1 & 0 \\ 0 & 11 \end{pmatrix}$  and  $M = \begin{pmatrix} 10 & 3i \\ -3i & 2 \end{pmatrix}$ . Similarly, transformation of  $M$  to  $\Lambda$  can be performed by


- (a)  $\frac{1}{\sqrt{10}} \begin{pmatrix} 1 & 3i \\ 3i & 1 \end{pmatrix}$                       (b)  $\frac{1}{\sqrt{9}} \begin{pmatrix} 1 & -3i \\ 3i & 11 \end{pmatrix}$   
 (c)  $\frac{1}{\sqrt{10}} \begin{pmatrix} 1 & 3i \\ -3i & 11 \end{pmatrix}$                       (d)  $\frac{1}{\sqrt{9}} \begin{pmatrix} 1 & 3i \\ -3i & 1 \end{pmatrix}$

-





- Q15. Consider the following circuit in steady state condition. Calculate the amount of charge stored in  $1\mu F$  and  $2\mu F$  capacitors respectively.



- Q16. If the mean square fluctuations in energy of a system in equilibrium at temperature  $T$  is proportional to  $T^\alpha$ , then the energy of the system is proportional to
- (a)  $T^{\alpha-2}$  (b)  $T^{\frac{\alpha}{2}}$  (c)  $T^{\alpha-1}$  (d)  $T^\alpha$
- Q17. Suppose the spin degrees of freedom of a 2-particle system can be described by a 21-dimensional Hilbert subspace. Which among the following could be the spin of one of the particles?
- (a)  $\frac{1}{2}$  (b) 3 (c)  $\frac{3}{2}$  (d) 2
- Q18. Water is poured at a rate of  $R \text{ m}^3/\text{hour}$  from the top into a cylindrical vessel of diameter  $D$ . The vessel has a small opening of area  $a(\sqrt{a} \ll D)$  at the bottom. What should be the minimum height of the vessel so that water does not overflow?
- (a)  $\infty$  (b)  $\frac{R^2}{2ga^2}$  (c)  $\frac{R^2}{2gaD^2}$  (d)  $\frac{8R^2}{\pi D^2 g^2}$
- Q19. Suppose that we toss two fair coins hundred times each. The probability that the same number of heads occur for both coins at the end of the experiment is
- (a)  $\left(\frac{1}{4}\right)^{100} \sum_{n=0}^{100} \binom{100}{n}$  (b)  $2\left(\frac{1}{4}\right)^{100} \sum_{n=0}^{100} \binom{100}{n}^2$
- (c)  $\frac{1}{2}\left(\frac{1}{4}\right)^{100} \sum_{n=0}^{100} \binom{100}{n}^2$  (d)  $\left(\frac{1}{4}\right)^{100} \sum_{n=0}^{100} \binom{100}{n}^2$
- Q20. What is the equation of the plane which is tangent to the surface  $xyz = 4$  at the point  $(1, 2, 2)$ ?
- (a)  $x + 2y + 4z = 12$  (b)  $4x + 2y + z = 12$
- (c)  $x + 4y + z = 20$  (d)  $2x + y + z = 6$
- Q21. If the ground state wavefunction of a particle moving in a one dimensional potential is proportional to  $\exp(-x^2/2)\cosh(\sqrt{2}x)$ , then the potential in suitable units such that  $\hbar = 1$ , is proportional to
- (a)  $x^2$  (b)  $x^2 - 2\sqrt{2}x \tanh(\sqrt{2}x)$
- (c)  $x^2 - 2\sqrt{2}x \tan(\sqrt{2}x)$  (d)  $x^2 - 2\sqrt{2}x \coth(\sqrt{2}x)$

Q22. A possible Lagrangian for a free particle is

- (a)  $L = \dot{q}^2 - q^2$  (b)  $L = \dot{q}^2 - q\dot{q}$   
(c)  $L = \dot{q}^2 - q$  (d)  $L = \dot{q}^2 - \frac{1}{q}$

Q23. A rod of mass  $m$  and length  $l$  is suspended from two massless vertical springs with a spring constants  $k_1$  and  $k_2$ . What is the Lagrangian for the system, if  $x_1$  and  $x_2$  be the displacements from equilibrium position of the two ends of the rod?

- (a)  $\frac{m}{8}(\dot{x}_1^2 + 2\dot{x}_1\dot{x}_2 + \dot{x}_2^2) - \frac{1}{2}k_1x_1^2 - \frac{1}{2}k_2x_2^2$   
(b)  $\frac{m}{2}(\dot{x}_1^2 + \dot{x}_1\dot{x}_2 + \dot{x}_2^2) - \frac{1}{4}(k_1 + k_2)(x_1^2 + x_2^2)$   
(c)  $\frac{m}{6}(\dot{x}_1^2 + \dot{x}_1\dot{x}_2 + \dot{x}_2^2) - \frac{1}{2}k_1x_1^2 - \frac{1}{2}k_2x_2^2$   
(d)  $\frac{m}{2}(\dot{x}_1^2 - 2\dot{x}_1\dot{x}_2 + \dot{x}_2^2) - \frac{1}{4}(k_1 - k_2)(x_1^2 + x_2^2)$

Q24. Two equal positive charges of magnitude  $+q$  separated by a distance  $d$  are surrounded by a uniformly charged thin spherical shell of radius  $2d$  bearing a total charge  $-2q$  and centred at the midpoint between the two positive charges. The net electric field at distance  $r$  from the midpoint ( $r \gg d$ ) is

- (a) zero (b) proportional to  $d$   
(c) proportional to  $1/r^3$  (d) proportional to  $1/r^4$

Q25. If the Hamiltonian of a classical particles is  $H = \frac{p_x^2 + p_y^2}{2m} + xy$ , then  $\langle x^2 + xy + y^2 \rangle$  at temperature  $T$  is equal to

- (a)  $k_B T$  (b)  $\frac{1}{2} k_B T$  (c)  $2k_B T$  (d)  $\frac{3}{2} k_B T$

Ans (a)

## Part-B: 3-Mark Questions

Q1. A solid, insulating sphere of radius  $1\text{ cm}$  has charge  $10^{-7}\text{ C}$  distributed uniformly over its volume. It is surrounded concentrically by a conducting thick spherical shell of inner radius  $2\text{ cm}$ , outer radius  $2.5\text{ cm}$  and is charged with  $-2 \times 10^{-7}\text{ C}$ . What is the electrostatic potential in Volts on the surface of the sphere?

Q2. A particle is described by the following Hamiltonian


$$\hat{H} = \frac{\hat{p}^2}{2m} + \frac{1}{2}m\omega^2\hat{x}^2 + \lambda\hat{x}^4$$

where the quartic term can be treated perturbatively. If  $\Delta E_0$  and  $\Delta E_1$  denote the energy correction of  $O(\lambda)$  to the ground state and the first excited state respectively, what is the fraction  $\Delta E_1 / \Delta E_0$ ?

Q3. A simple pendulum has a bob of mass  $1\text{ kg}$  and charge  $1\text{ Coulomb}$ . It is suspended by a massless string of length  $13\text{ m}$ . The time period of small oscillations of this pendulum is  $T_0$ . If an electric field  $\vec{E} = 100\hat{x}\text{ V/m}$  is applied, the time period becomes  $T$ . What is the value of  $(T_0/T)^4$ ?

Q4. Let a particle of mass  $1 \times 10^{-9}\text{ kg}$ , constrained to have one dimensional motion, be initially at the origin ( $x = 0\text{ m}$ ). The particle is in equilibrium with a thermal bath ( $k_B T = 10^{-8}\text{ J}$ ). What is  $\langle x^2 \rangle$  of the particle after a time  $t = 5\text{ s}$ ?

Q5. For the circuit shown below, what is the ratio  $\frac{I_1}{I_2}$ ?





- Q6. A ball of mass  $0.1\text{ kg}$  and density  $2000\text{ kg/m}^3$  is suspended by a massless string of length  $0.5\text{ m}$  under water having density  $1000\text{ kg/m}^3$ . The ball experiences a drag force,  $\vec{F}_d = -0.2(\vec{v}_b - \vec{v}_w)$ , where  $\vec{v}_b$  and  $\vec{v}_w$  are the velocities of the ball and water respectively. What will be the frequency of small oscillations for the motion of pendulum, if the water is at rest?
- Q7. Suppose that the number of microstates available to a system of  $N$  particles depends on  $N$  and the combined variable  $UV^2$ , where  $U$  is the internal energy and  $V$  is the volume of the system. The system initially has volume  $2\text{ m}^3$  and energy  $200\text{ J}$ . It undergoes an isentropic expansion to volume  $4\text{ m}^3$ . What is the final pressure of the system in SI units?
- Q8. The temperature in a rectangular plate bounded by the lines,  $x=0, y=0, x=3$  and  $y=5$  is  $T = xy^2 - x^2y + 100$ . What is the maximum temperature difference between two points on the plate?
- Q9. A sphere of inner radius  $1\text{ cm}$  and outer radius  $2\text{ cm}$ , centered at origin has a volume charge density  $\rho_0 = \frac{K}{4\pi r}$ , where  $K$  is a nonzero constant and  $r$  is the radial distance. A point charge of magnitude  $10^{-3}\text{ C}$  is placed at the origin. For what value of  $K$  in units of  $\text{C/m}^2$  the electric field inside shell is constant?
- Q10. If  $\hat{x}(t)$  be the position operator at a time  $t$  in the Heisenberg picture for a particle described by the Hamiltonian,  $\hat{H} = \frac{\hat{p}^2}{2m} + \frac{1}{2}m\omega^2\hat{x}^2$  what is  $e^{i\omega t} \langle 0 | \hat{x}(t) \hat{x}(0) | 0 \rangle$  in units of  $\frac{\hbar}{2m\omega}$  where  $|0\rangle$  is the ground state?

### Part-C: 3-Mark Questions

- Q1. Consider a grounded conducting plane which is infinitely extended perpendicular to the  $y$ -axis at  $y=0$ . If an infinite line of charge per unit length  $\lambda$  runs parallel to  $x$ -axis at  $y=d$ , then surface charge density on the conducting plane is
- (a)  $\frac{-\lambda d}{(x^2 + d^2 + z^2)}$                       (b)  $\frac{-\lambda d}{(x^2 + d^2 + z^2)}$
- (c)  $\frac{-\lambda d}{\pi(x^2 + d^2 + z^2)}$                       (d)  $\frac{-\lambda d}{2\pi(x^2 + d^2 + z^2)}$


Q2. A system of particles on  $N$  lattice sites is in equilibrium at temperature  $T$  and chemical potential  $\mu$ . Multiple occupancy of the sites is forbidden. The binding energy of a particle at each site is  $-\epsilon$ . The probability of no site being occupied is

- (a)  $\frac{1 - e^{\beta(\mu + \epsilon)}}{1 - e^{(N+1)\beta(\mu + \epsilon)}}$       (b)  $\frac{1}{[1 + e^{\beta(\mu + \epsilon)}]^N}$
- (c)  $\frac{1}{[1 + e^{-\beta(\mu + \epsilon)}]^N}$       (d)  $\frac{1 - e^{\beta(\mu + \epsilon)}}{1 - e^{-(N+1)\beta(\mu + \epsilon)}}$

Q3. The integral  $I = \int_1^\infty \frac{\sqrt{x-1}}{(1+x)^2} dx$  is

- (a)  $\frac{\pi}{\sqrt{2}}$       (b)  $\frac{\pi}{2\sqrt{2}}$       (c)  $\frac{\sqrt{\pi}}{2}$       (d)  $\sqrt{\frac{\pi}{2}}$

Q4. For an electric field  $\vec{E} = k\sqrt{x}\hat{x}$  where  $k$  is a non-zero constant, total charge enclosed by the cube as shown below is



- (a) 0      (b)  $k \epsilon_0 l^{5/2} (\sqrt{3} - 1)$
- (c)  $k \epsilon_0 l^{5/2} (\sqrt{5} - 1)$       (d)  $k \epsilon_0 l^{5/2} (\sqrt{2} - 1)$


Q5. Consider a point particle  $A$  of mass  $m_A$  colliding elastically with another point particle  $B$  of mass  $m_B$  at rest, where  $\frac{m_B}{m_A} = \gamma$ . After collision, the ratio of the kinetic energy of particle  $B$  to the initial kinetic energy of particle  $A$  is given by

- (a)  $\frac{4}{\gamma + 2 + \frac{1}{\gamma}}$       (b)  $\frac{2}{\gamma + \frac{1}{\gamma}}$
- (c)  $\frac{2}{\gamma + 2 - \frac{1}{\gamma}}$       (d)  $\frac{1}{\gamma}$

Q6. Two classical particles are distributed among  $N(>2)$  sites on a ring. Each site can accommodate only one particle. If two particles occupy two nearest neighbour sites, then the energy of the system is increased by  $\epsilon$ . The average energy of the system at temperature  $T$  is

- (a)  $\frac{2\epsilon e^{-\beta\epsilon}}{(N-3)+2e^{-\beta\epsilon}}$                       (b)  $\frac{2N\epsilon e^{-\beta\epsilon}}{(N-3)+2e^{-\beta\epsilon}}$
- (c)  $\frac{\epsilon}{N}$     (d)  $\frac{2\epsilon e^{-\beta\epsilon}}{(N-2)+2e^{-\beta\epsilon}}$

Q7. Consider a 741 operational amplifier circuit as shown below, where  $V_{CC} = V_{EE} = +15V$  and  $R = 2.2k\Omega$ . If  $v_i = 2mV$ , what is the value of  $v_o$  with respect to the ground?



- (a)  $-1mV$                       (b)  $-2mV$                       (c)  $-3mV$                       (d)  $-4mV$

Q8. The Fourier transform of the function  $\frac{1}{x^4 + 3x^2 + 2}$  up to proportionality constant is

- (a)  $\sqrt{2} \exp(-k^2) - \exp(-2k^2)$                       (b)  $\sqrt{2} \exp(-|k|) - \exp(-\sqrt{2}|k|)$
- (c)  $\sqrt{2} \exp(-\sqrt{|k|}) - \exp(-\sqrt{2}|k|)$                       (d)  $\sqrt{2} \exp(-\sqrt{2}k^2) - \exp(-2k^2)$

Q9. If  $\rho = \frac{\left[ I + \frac{1}{\sqrt{3}}(\sigma_x + \sigma_y + \sigma_z) \right]}{2}$ , where  $\sigma$ 's are the Pauli matrices and  $I$  is the identity matrix, then the trace of  $\sigma^{2017}$  is


- (a)  $2^{2017}$                       (b)  $2^{-2017}$                       (c) 1                      (d)  $\frac{1}{2}$

Q10. A cylinder at temperature  $T = 0$  is separated into two compartments  $A$  and  $B$  by a free sliding piston. Compartments  $A$  and  $B$  are filled by Fermi gases made of spin  $1/2$  and  $3/2$  particles respectively. If particles in both the compartments have same mass, the ratio of equilibrium density of the gas in compartment  $A$  to that of gas in compartment  $B$  is

- (a) 1                      (b)  $\frac{1}{3^{2/5}}$                       (c)  $\frac{1}{2^{2/5}}$                       (d)  $\frac{1}{2^{2/3}}$

Q11. What is the DC base current (approximated to nearest integer value in  $\mu A$ ) for the following  $n - p - n$  silicon transistor circuit,

given  $R_1 = 75 \Omega$ ,  $R_2 = 4.0 k\Omega$ ,  $R_3 = 2.1 k\Omega$ ,  $R_4 = 2.6 k\Omega$ ,  $R_5 = 6.0 k\Omega$ ,



- (a) 20                      (b) 24                      (c) 16                      (d) 32

Q12. Consider a particle confined by a potential  $V(x) = k|x|$ , where  $k$  is a positive constant. The spectrum  $E_n$  of the system, within the WKB approximation is proportional to

- (a)  $\left(n + \frac{1}{2}\right)^{3/2}$                       (b)  $\left(n + \frac{1}{2}\right)^{2/3}$                       (c)  $\left(n + \frac{1}{2}\right)^{1/2}$                       (d)  $\left(n + \frac{1}{2}\right)^{4/3}$

Q13. Consider the Hamiltonian

$$H(t) = \alpha \begin{pmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 3 \end{pmatrix} + \beta t \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 1 & 0 & -2 \end{pmatrix}$$

The time dependent function  $\beta(t) = \alpha$  for  $t \leq 0$  and zero for  $t > 0$ . Find  $\left| \langle \Psi(t < 0) | \Psi(t > 0) \rangle \right|^2$ , where  $|\Psi(t < 0)\rangle$  is the normalised ground state of the system at a time  $t < 0$  and  $|\Psi(t > 0)\rangle$  is the state of the system at  $t > 0$ .

- (a)  $\frac{1}{2}(1 + \cos(2\alpha t))$                       (b)  $\frac{1}{2}(1 + \cos(\alpha t))$

(c)  $\frac{1}{2}(1 + \sin(2\alpha t))$

(d)  $\frac{1}{2}(1 + \sin(\alpha t))$

Q14. The function  $f(x) = \cosh x$  which exists in the range  $-\pi \leq x \leq \pi$  is periodically repeated between  $x = (2m-1)\pi$  and  $(2m+1)\pi$ , where  $m = -\infty$  to  $\infty$ . Using Fourier series, indicate the correct relation at  $x = 0$

(a)  $\sum_{n=-\infty}^{\infty} \frac{(-1)^n}{1-n^2} = \frac{1}{2} \left( \frac{\pi}{\cosh \pi} - 1 \right)$

(b)  $\sum_{n=-\infty}^{\infty} \frac{(-1)^n}{1-n^2} = 2 \frac{\pi}{\cosh \pi}$

(c)  $\sum_{n=-\infty}^{\infty} \frac{(-1)^{-n}}{1+n^2} = 2 \frac{\pi}{\sinh \pi}$

(d)  $\sum_{n=1}^{\infty} \frac{(-1)^n}{1+n^2} = \frac{1}{2} \left( \frac{\pi}{\sinh \pi} - 1 \right)$

Q15. A toy car is made from a rectangular block of mass  $M$  and four disk wheels of mass  $m$  and radii  $r$ . The car is attached to a vertical wall by a massless horizontal spring with spring constant  $k$  and constrained to move perpendicular to the wall. The coefficient of static friction between the wheel of the car and the floor is  $\mu$ . The maximum amplitude of oscillations of the car above which the wheels start slipping is

(a)  $\frac{\mu g (M + 2m)(M + 4m)}{mk}$

(b)  $\frac{\mu g (M^2 - m^2)}{Mk}$

(c)  $\frac{\mu g (M + m)^2}{2mk}$

(d)  $\frac{\mu g (M + 4m)(M + 6m)}{2mk}$