


## NET-JRF (December 2018)

### PART- A

- Q1. A rectangular photo frame of size  $30\text{ cm} \times 40\text{ cm}$  has a photograph mounted at the centre leaving a  $5\text{ cm}$  border all around. The area of the border is
- (a)  $600\text{ cm}^2$                       (b)  $350\text{ cm}^2$                       (c)  $400\text{ cm}^2$                       (d)  $700\text{ cm}^2$
- Q2. At a birthday party, every child gets 2 chocolates, every mother gets 1 chocolate, while no father gets a chocolate. In total 69 persons get 70 chocolates. If the number of children is half of the number of mothers and fathers put together, then how many fathers are there?
- (a) 22                                      (b) 23                                      (c) 24                                      (d) 69
- Q3. What is the value of  $1^2 - 2^2 + 3^2 - 4^2 + 5^2 - \dots + 17^2 - 18^2 + 19^2$ ?
- (a)  $-5$                                       (b) 12                                      (c) 95                                      (d) 190
- Q4. The curves of  $y = 2x^2$  and  $y = 4x$  intersect each other at
- (a) only one point                                      (b) exactly two points  
(c) more than two points                                      (d) no point at all
- Q5. The diameters of the pinholes of two otherwise identical cameras  $A$  and  $B$  are  $500\ \mu\text{m}$  and  $200\ \mu\text{m}$ , respectively. Then the image in camera  $A$  will be
- (a) sharper than in  $B$                                       (b) darker than in  $B$   
(c) less sharp and brighter than in  $B$                                       (d) sharper and brighter than in  $B$
- Q6. If  $D = ABC + BCA + CAB$ , where  $A$ ,  $B$  and  $C$  are decimal digits, then  $D$  is divisible by
- (a) 37 and 29                                      (b) 37 but not 29  
(c) 29 but not 37                                      (d) neither 29 nor 37
- Q7. For the following set of observed values
- $$\{60, 65, 65, 70, 70, 70, 70, 82, 85, 90, 95, 95, 100, 160, 160\}$$
- which of the statements is true?
- (a) mode < median < mean                                      (b) mode < mean < median  
(c) mean < median < mode                                      (d) median < mode < mean

- Q8. A circular running track has six lanes, each  $1\text{ m}$  wide. How far ahead (in meters) should the runner in the outermost lane start from, so as to cover the same distance in one lap as the runner in the innermost lane?
- (a)  $6\pi$                       (b)  $10\pi$                       (c)  $12\pi$                       (d)  $36\pi$
- Q9. In an examination 100 questions of 1 mark each are given. After the examination, 20 questions are deleted from evaluation, leaving 80 questions with a total of 100 marks. Student  $A$  had answered 4 of the deleted questions correctly and got 40 marks, whereas student  $B$  had answered 10 of the deleted questions correctly and got 35 marks. In this situation
- (a)  $A$  and  $B$  were equally benefited                      (b)  $A$  and  $B$  lost equally  
(c)  $B$  lost more than  $A$                       (d)  $A$  lost more than  $B$
- Q10. A tourist drives  $20\text{ km}$  towards east, turns right and drives  $6\text{ km}$ , then drives  $6\text{ km}$  towards west. He then turns to his left and drives  $4\text{ km}$  and finally turns right and drives  $14\text{ km}$ . Where is he from his starting point?
- (a)  $6\text{ km}$  towards east                      (b)  $20\text{ km}$  towards west  
(c)  $14\text{ km}$  towards north                      (d)  $10\text{ km}$  towards south
- Q11. If 'SELDOON' means 'NOODLES' then what does 'SPUOS' mean?
- (a) SALAD                      (b) SOUPS                      (c) RASAM                      (d) ONION
- Q12. An ideal pendulum oscillates with angular amplitude of  $30^\circ$  from the vertical. If it is observed at a random instant of time, its angular deviation from the vertical is most likely to be
- (a)  $0^\circ$                       (b)  $\pm 10^\circ$                       (c)  $\pm 20^\circ$                       (d)  $\pm 30^\circ$
- Q13. In the context of tiling a plane surface, which of the following polygons is the odd one out?
- (a) Equilateral triangle                      (b) Square  
(c) Regular pentagon                      (d) Regular hexagon

Q14. Scatter plots for pairs of observations on the variables  $x$  and  $y$  in samples  $A$  and  $B$  are shown in the figure.




Which of the following is suggested by the plots?

- (a) Correlation between  $x$  and  $y$  is stronger in  $A$  than in  $B$
  - (b) Correlation between  $x$  and  $y$  is absent in  $B$
  - (c) Correlation between  $x$  and  $y$  is weaker in  $A$  than in  $B$
  - (d)  $y$  and  $x$  have a cause - effect relationship in  $A$  but not in  $B$
- Q15. Two solutions  $X$  and  $Y$  containing ingredients  $A, B$  and  $C$  in proportions  $a:b:c$  and  $c:b:a$ , respectively, are mixed. For the resultant mixture to have  $A, B$  and  $C$  in equal proportion, it is necessary that


- (a)  $b = \frac{c-a}{2}$
- (b)  $c = \frac{a+b}{2}$
- (c)  $c = \frac{a-b}{2}$
- (d)  $b = \frac{c+a}{2}$

Q16. Find the missing figure in the following sequence.



- (a)
- (b)
- (c)
- (d)

Q17. In triangle  $ABC$ ,  $AB = 11$ ,  $BC = 61$ ,  $AC = 60$ , and  $O$  is the mid-point of  $BC$ . Then  $AO$  is



- (a) 18.5                      (b) 24.0                      (c) 30.5                      (d) 36.0

Q18. Areas of three parts of a rectangle are given in unit of  $cm^2$ . What is the total area of the rectangle?

3	9
6	

- (a) 18                      (b) 24                      (c) 36                      (d) 108

Q19. A student is free to choose only Chemistry, only Biology or both. If out of 32 students, Chemistry has been chosen by 16 and Biology by 25, then how many students have chosen Biology but not Chemistry?

- (a) 9                      (b) 16                      (c) 25                      (d) 7

Q20. The lift (upward force due to air) generated by the wings and engines of an aircraft is

- (a) positive (upwards) while landing and negative (downwards) while taking off.  
(b) negative (downwards) while landing and positive (upwards) while taking off  
(c) negative (downwards) while landing as well as while taking off  
(d) positive (upwards) while landing as well as while taking off

## PART B

Q21. One of the eigenvalues of the matrix  $e^A$  is  $e^a$ , where  $A = \begin{pmatrix} a & 0 & 0 \\ 0 & 0 & a \\ 0 & a & 0 \end{pmatrix}$ . The product of the other

two eigenvalues of  $e^A$  is

- (a)  $e^{2a}$                       (b)  $e^{-a}$                       (c)  $e^{-2a}$                       (d) 1

Q22. The polynomial  $f(x) = 1 + 5x + 3x^2$  is written as linear combination of the Legendre polynomials

$\left( p_0(x) = 1, p_1(x), p_2(x) = \frac{1}{2}(3x^2 - 1) \right)$  as  $f(x) = \sum_n c_n p_n(x)$ . The value of  $c_0$  is

- (a)  $\frac{1}{4}$                       (b)  $\frac{1}{2}$                       (c) 2                      (d) 4


Q23. The value of the integral  $\oint_C \frac{dz \tanh 2z}{z \sin \pi z}$ , where  $C$  is a circle of radius  $\frac{\pi}{2}$ , traversed counter-clockwise, with centre at  $z = 0$ , is

- (a) 4                      (b)  $4i$                       (c)  $2i$                       (d) 0

Q24. 1(a)  $v_0 e^{-\frac{\gamma v_0 t}{m}}$                       (b)  $\frac{v_0}{1 + \ln\left(1 + \frac{\gamma v_0 t}{m}\right)}$

(c)  $\frac{m v_0}{m + \gamma v_0 t}$                       (d)  $\frac{2 v_0}{1 + e^{\frac{\gamma v_0 t}{m}}}$

Q25. The integral  $I = \int_C e^z dz$  is evaluated from the point  $(-1, 0)$  to  $(1, 0)$  along the contour  $C$ , which is an arc of the parabola  $y = x^2 - 1$ , as shown in the figure.



The value of  $I$  is

- (a) 0                      (b)  $2 \sinh 1$                       (c)  $e^{2i} \sinh 1$                       (d)  $e + e^{-1}$

Q26. In terms of arbitrary constants  $A$  and  $B$ , the general solution to the differential equation

$$x^2 \frac{d^2 y}{dx^2} + 5x \frac{dy}{dx} + 3y = 0 \text{ is}$$

(a)  $y = \frac{A}{x} + BX^3$

(b)  $y = Ax + \frac{B}{x^3}$

(c)  $y = Ax + Bx^3$

(d)  $y = \frac{A}{x} + \frac{B}{x^3}$

Q27. In the attractive Kepler problem described by the central potential  $V(r) = \frac{-k}{r}$  (where  $k$  is a positive constant), a particle of mass  $m$  with a non-zero angular momentum can never reach the centre due to the centrifugal barrier. If we modify the potential to

$$V(r) = -\frac{k}{r} - \frac{\beta}{r^3}$$

one finds that there is a critical value of the angular momentum  $\ell_c$  below which there is no centrifugal barrier. This value of  $\ell_c$  is

(a)  $[12km^2\beta]^{1/2}$

(b)  $[12km^2\beta]^{-1/2}$

(c)  $[12km^2\beta]^{1/4}$

(d)  $[12km^2\beta]^{-1/4}$

Q28. The time period of a particle of mass  $m$ , undergoing small oscillations around  $x = 0$ , in the potential  $V = V_0 \cosh\left(\frac{x}{L}\right)$ , is

(a)  $\pi \sqrt{\frac{mL^2}{v_0}}$

(b)  $2\pi \sqrt{\frac{mL^2}{2v_0}}$

(c)  $2\pi \sqrt{\frac{mL^2}{v_0}}$

(d)  $2\pi \sqrt{\frac{2mL^2}{v_0}}$

Q29. Consider the decay  $A \rightarrow B + C$  of a relativistic spin- $\frac{1}{2}$  particle  $A$ . Which of the following statements is true in the rest frame of the particle  $A$ ?

(a) The spin of both  $B$  and  $C$  may be  $\frac{1}{2}$

(b) The sum of the masses of  $B$  and  $C$  is greater than the mass of  $A$

(c) The energy of  $B$  is uniquely determined by the masses of the particles

(d) The spin of both  $B$  and  $C$  may be integral

- Q30. Two current-carrying circular loops, each of radius  $R$ , are placed perpendicular to each other, as shown in the figure.

The loop in the  $xy$ -plane carries a current  $I_0$  while that in the  $xz$ -plane carries a current  $2I_0$ .

The resulting magnetic field  $\vec{B}$  at the origin is

- (a)  $\frac{\mu_0 I_0}{2R} [2\hat{j} + \hat{k}]$                       (b)  $\frac{\mu_0 I_0}{2R} [2\hat{j} - \hat{k}]$   
(c)  $\frac{\mu_0 I_0}{2R} [-2\hat{j} + \hat{k}]$                       (d)  $\frac{\mu_0 I_0}{2R} [-2\hat{j} - \hat{k}]$

- Q31. An electric dipole of dipole moment  $\vec{P} = qb\hat{i}$  is placed at origin in the vicinity of two charges  $+q$  and  $-q$  at  $(L, b)$  and  $(L, -b)$ , respectively, as shown in the figure below.

The electrostatic potential at the point  $\left(\frac{L}{2}, 0\right)$  is

- (a)  $\frac{qb}{\pi \epsilon_0} \left( \frac{1}{L^2} + \frac{2}{L^2 + 4b^2} \right)$                       (b)  $\frac{4qbL}{\pi \epsilon_0 [L^2 + 4b^2]^{3/2}}$   
(c)  $\frac{qb}{\pi \epsilon_0 L^2}$                       (d)  $\frac{3qb}{\pi \epsilon_0 L^2}$

- Q32. A monochromatic and linearly polarized light is used in a Young's double slit experiment. A linear polarizer, whose pass axis is at an angle  $45^\circ$  to the polarization of the incident wave, is placed in front of one of the slits. If  $I_{\max}$  and  $I_{\min}$ , respectively, denote the maximum and minimum intensities of the interference pattern on the screen, the visibility, defined as the

ratio  $\frac{I_{\max} - I_{\min}}{I_{\max} + I_{\min}}$ , is

- (a)  $\frac{\sqrt{2}}{3}$                       (b)  $\frac{2}{3}$                       (c)  $\frac{2\sqrt{2}}{3}$                       (d)  $\sqrt{\frac{2}{3}}$

Q33. An electromagnetic wave propagates in a nonmagnetic medium with relative permittivity  $\epsilon = 4$ .

The magnetic field for this wave is

$$\vec{H}(x, y) = \hat{k}H_0 \cos(\omega t - \alpha x - \alpha\sqrt{3}y)$$

where  $H_0$  is a constant. The corresponding electric field  $\vec{E}(x, y)$  is

(a)  $\frac{1}{4} \mu_0 H_0 c (-\sqrt{3}\hat{i} + \hat{j}) \cos(\omega t - \alpha x - \alpha\sqrt{3}y)$

(b)  $\frac{1}{4} \mu_0 H_0 c (\sqrt{3}\hat{i} + \hat{j}) \cos(\omega t - \alpha x - \alpha\sqrt{3}y)$

(c)  $\frac{1}{4} \mu_0 H_0 c (\sqrt{3}\hat{i} - \hat{j}) \cos(\omega t - \alpha x - \alpha\sqrt{3}y)$

(d)  $\frac{1}{4} \mu_0 H_0 c (-\sqrt{3}\hat{i} - \hat{j}) \cos(\omega t - \alpha x - \alpha\sqrt{3}y)$

Q34. The ground state energy of an anisotropic harmonic oscillator described by the potential

$$V(x, y, z) = \frac{1}{2} m\omega^2 x^2 + 2m\omega^2 y^2 + 8m\omega^2 z^2 \text{ (in units of } \hbar\omega \text{ ) is}$$

(a)  $\frac{5}{2}$

(b)  $\frac{7}{2}$

(c)  $\frac{3}{2}$

(d)  $\frac{1}{2}$

Q35. The product  $\Delta x \Delta p$  of uncertainties in the position and momentum of a simple harmonic

oscillator of mass  $m$  and angular frequency  $\omega$  in the ground state  $|0\rangle$ , is  $\frac{\hbar}{2}$ . The value of the

product  $\Delta x \Delta p$  in the state,  $e^{-i\hat{p}\ell/\hbar} |0\rangle$  (where  $\ell$  is a constant and  $\hat{p}$  is the momentum operator) is

(a)  $\frac{\hbar}{2} \sqrt{\frac{m\omega\ell^2}{\hbar}}$

(b)  $\hbar$

(c)  $\frac{\hbar}{2}$

(d)  $\frac{\hbar^2}{m\omega\ell^2}$

Q36. Let the wavefunction of the electron in a hydrogen atom be

$$\psi(\vec{r}) = \frac{1}{\sqrt{6}} \phi_{200}(\vec{r}) + \sqrt{\frac{2}{3}} \phi_{21-1}(\vec{r}) - \frac{1}{\sqrt{6}} \phi_{100}(\vec{r})$$

where  $\phi_{nlm}(\vec{r})$  are the eigenstates of the Hamiltonian in the standard notation. The expectation value of the energy in this state is

(a)  $-10.8 \text{ eV}$

(b)  $-6.2 \text{ eV}$

(c)  $-9.5 \text{ eV}$

(d)  $-5.1 \text{ eV}$




- Q37. Three identical spin  $\frac{1}{2}$  particles of mass  $m$  are confined to a one-dimensional box of length  $L$ , but are otherwise free. Assuming that they are non-interacting, the energies of the lowest two energy eigen states, in units of  $\frac{\pi^2 \hbar^2}{2mL^2}$ , are
- (a) 3 and 6                      (b) 6 and 9                      (c) 6 and 11                      (d) 3 and 9
- Q38. The heat capacity  $C_V$  at constant volume of a metal, as a function of temperature, is  $\alpha T + \beta T^3$ , where  $\alpha$  and  $\beta$  are constant. The temperature dependence of the entropy at constant volume is
- (a)  $\alpha T + \frac{1}{3} \beta T^3$                       (b)  $\alpha T + \beta T^3$   
(c)  $\frac{1}{2} \alpha T + \frac{1}{3} \beta T^3$                       (d)  $\frac{1}{2} \alpha T + \frac{1}{4} \beta T^3$
- Q39. The rotational energy levels of a molecule are  $E_\ell = \frac{\hbar^2}{2I_0} \ell(\ell+1)$ , where  $\ell = 0, 1, 2, \dots$  and  $I_0$  is its moment of inertia. The contribution of the rotational motion to the Helmholtz free energy per molecule, at low temperatures in a dilute gas of these molecules, is approximately
- (a)  $-k_B T \left( 1 + \frac{\hbar^2}{I_0 k_B T} \right)$                       (b)  $-k_B T e^{-\frac{\hbar^2}{I_0 k_B T}}$   
(c)  $-k_B T$                       (d)  $-3k_B T e^{-\frac{\hbar^2}{I_0 k_B T}}$
- Q40. The vibrational motion of a diatomic molecule may be considered to be that of a simple harmonic oscillator with angular frequency  $\omega$ . If a gas of these molecules is at temperature  $T$ , what is the probability that a randomly picked molecule will be found in its lowest vibrational state?
- (a)  $1 - e^{-\frac{\hbar\omega}{k_B T}}$                       (b)  $e^{-\frac{\hbar\omega}{2k_B T}}$   
(c)  $\tanh\left(\frac{\hbar\omega}{k_B T}\right)$                       (d)  $\frac{1}{2} \operatorname{cosec} h\left(\frac{\hbar\omega}{2k_B T}\right)$

Q41. Consider an ideal Fermi gas in a grand canonical ensemble at a constant chemical potential. The variance of the occupation number of the single particle energy level with mean occupation number  $\bar{n}$  is

- (a)  $\bar{n}(1-\bar{n})$       (b)  $\sqrt{\bar{n}}$       (c)  $\bar{n}$       (d)  $\frac{1}{\sqrt{\bar{n}}}$

Q42. Consider the following circuit, consisting of an  $RS$  flip-flop and two AND gates.



Which of the following connections will allow the entire circuit to act as a  $JK$  flip-flop?

- (a) connect  $Q$  to pin 1 and  $\bar{Q}$  to pin 2  
 (b) connect  $Q$  to pin 2 and  $\bar{Q}$  to pin 1  
 (c) connect  $Q$  to  $K$  input and  $\bar{Q}$  to  $J$  input  
 (d) connect  $Q$  to  $J$  input and  $\bar{Q}$  to  $K$  input


Q43. The truth table below gives the value  $Y(A, B, C)$  where  $A, B$  and  $C$  are binary variables.

$A$	$B$	$C$	$Y$
0	0	0	1
0	0	1	0
0	1	0	0
0	1	1	1
1	0	0	1
1	1	0	0
1	1	1	1


The output  $Y$  can be represented by

- (a)  $Y = \bar{A}\bar{B}C + \bar{A}B\bar{C} + A\bar{B}C + A\bar{B}\bar{C}$   
 (b)  $Y = \bar{A}\bar{B}\bar{C} + \bar{A}BC + A\bar{B}\bar{C} + ABC$   
 (c)  $Y = \bar{A}\bar{B}C + \bar{A}BC + A\bar{B}C + ABC$   
 (d)  $Y = \bar{A}\bar{B}\bar{C} + \bar{A}B\bar{C} + A\bar{B}\bar{C} + ABC$


Q44. A sinusoidal signal is an input to the following circuit







Which of the following graphs best describes the output wave function?



Q45. A sinusoidal voltage having a peak value of  $V_p$  is an input to the following circuit, in which the DC voltage is  $V_b$



Assuming an ideal diode which of the following best describes the output waveform?



### PART C

Q46. The Green's function  $G(x, x')$  for the equation  $\frac{d^2 y(x)}{dx^2} = f(x)$ , with the boundary values  $y(0) = 0$  and  $y(1) = 0$ , is

(a)  $G(x, x') = \begin{cases} \frac{1}{2}x(1-x), & 0 < x < x' < 1 \\ \frac{1}{2}x'(1-x) & 0 < x' < x < 1 \end{cases}$

(b)  $G(x, x') = \begin{cases} x(x'-1), & 0 < x < x' < 1 \\ x'(1-x) & 0 < x' < x < 1 \end{cases}$

(c)  $G(x, x') = \begin{cases} -\frac{1}{2}x(1-x'), & 0 < x < x' < 1 \\ \frac{1}{2}x'(1-x) & 0 < x' < x < 1 \end{cases}$

(d)  $G(x, x') = \begin{cases} x(x'-1), & 0 < x < x' < 1 \\ x'(x-1) & 0 < x' < x < 1 \end{cases}$

Q47. A  $4 \times 4$  complex matrix  $A$  satisfies the relation  $A^\dagger A = 4I$ , where  $I$  is the  $4 \times 4$  identity matrix.


The number of independent real parameters of  $A$  is

- (a) 32                      (b) 10                      (c) 12                      (d) 16


Q48. The contour  $C$  of the following integral


$$\oint_C dz \frac{\sqrt{(z-1)(z-3)}}{(z^2-25)^3}$$

is the complex  $z$  plane is shown in the figure below.



This integral is equivalent to an integral along the contours





Q49. The value of the integral  $\int_0^1 x^2 dx$ , evaluated using the trapezoidal rule with a step size of 0.2, is  
 (a) 0.30                      (b) 0.39                      (c) 0.34                      (d) 0.27

Q50. The motion of a particle in one dimension is described by the Lagrangian  $L = \frac{1}{2} \left( \left( \frac{dx}{dt} \right)^2 - x^2 \right)$  in suitable units. The value of the action along the classical path from  $x = 0$  at  $t = 0$  to  $x = x_0$  at  $t = t_0$ , is

- (a)  $\frac{x_0^2}{2 \sin^2 t_0}$                       (b)  $\frac{1}{2} x_0^2 \tan t_0$                       (c)  $\frac{1}{2} x_0^2 \cot t_0$                       (d)  $\frac{x_0^2}{2 \cos^2 t_0}$

Q51. The Hamiltonian of a classical one-dimensional harmonic oscillator is

$H = \frac{1}{2} (p^2 + x^2)$ , in suitable units. The total time derivative of the dynamical variable

$(p + \sqrt{2}x)$  is

- (a)  $\sqrt{2}p - x$                       (b)  $p - \sqrt{2}x$                       (c)  $p + \sqrt{2}x$                       (d)  $x + \sqrt{2}p$

Q52. A relativistic particle of mass  $m$  and charge  $e$  is moving in a uniform electric field of strength  $\epsilon$ .

Starting from rest at  $t = 0$ , how much time will it take to reach the speed  $\frac{c}{2}$ ?

- (a)  $\frac{1}{\sqrt{3}} \frac{mc}{e\epsilon}$                       (b)  $\frac{mc}{e\epsilon}$                       (c)  $\sqrt{2} \frac{mc}{e\epsilon}$                       (d)  $\sqrt{\frac{3}{2}} \frac{mc}{e\epsilon}$

Q53. In an inertial frame uniform electric and magnetic field  $\vec{E}$  and  $\vec{B}$  are perpendicular to each other and satisfy  $|\vec{E}|^2 - |\vec{B}|^2 = 29$  (in suitable units). In another inertial frame, which moves at a constant velocity with respect to the first frame, the magnetic field is  $2\sqrt{5}\hat{k}$ . In the second frame, an electric field consistent with the previous observations is

- (a)  $\frac{7}{\sqrt{2}}(\hat{i} + \hat{j})$                       (b)  $7(\hat{i} + \hat{k})$                       (c)  $\frac{7}{\sqrt{2}}(\hat{i} + \hat{k})$                       (d)  $7(\hat{i} + \hat{j})$

- Q54. Electromagnetic wave of angular frequency  $\omega$  is propagating in a medium in which, over a band of frequencies the refractive index is  $n(\omega) \approx 1 - \left(\frac{\omega}{\omega_0}\right)^2$ , where  $\omega_0$  is a constant. The ratio  $\frac{v_g}{v_p}$  of the group velocity to the phase velocity at  $\omega = \frac{\omega_0}{2}$  is
- (a) 3                      (b)  $\frac{1}{4}$                       (c)  $\frac{2}{3}$                       (d) 2
- Q55. A rotating spherical shell of uniform surface charge and mass density has total mass  $M$  and charge  $Q$ . If its angular momentum is  $L$  and magnetic moment is  $\mu$ , then the ratio  $\frac{\mu}{L}$  is
- (a)  $\frac{Q}{3M}$                       (b)  $\frac{2Q}{3M}$                       (c)  $\frac{Q}{2M}$                       (d)  $\frac{3Q}{4M}$
- Q56. Consider the operator  $A_x = L_y p_z - L_z p_y$ , where  $L_i$  and  $p_i$  denote, respectively, the components of the angular momentum and momentum operators. The commutator  $[A_x, x]$ , where  $x$  is the  $x$ -component of the position operator, is
- (a)  $-i\hbar(zp_z + yp_y)$                       (b)  $-i\hbar(zp_z - yp_y)$   
(c)  $i\hbar(zp_z + yp_y)$                       (d)  $i\hbar(zp_z - yp_y)$
- Q57. A one-dimensional system is described by the Hamiltonian  $H = \frac{p^2}{2m} + \lambda|x|$  (where  $\lambda > 0$ ). The ground state energy varies as a function of  $\lambda$  as
- (a)  $\lambda^{5/3}$                       (b)  $\lambda^{2/3}$                       (c)  $\lambda^{4/3}$                       (d)  $\lambda^{1/3}$
- Q58. If the position of the electron in the ground state of a Hydrogen atom is measured, the probability that it will be found at a distance  $r \geq a_0$  ( $a_0$  being Bohr radius) is nearest to
- (a) 0.91                      (b) 0.66                      (c) 0.32                      (d) 0.13
- Q59. A system of spin  $\frac{1}{2}$  particles is prepared to be in the eigenstate of  $\sigma_z$  with eigenvalue +1. The system is rotated by an angle of  $60^\circ$  about the  $x$ -axis. After the rotation, the fraction of the particles that will be measured to be in the eigenstate of  $\sigma_z$  with eigenvalue +1 is
- (a)  $\frac{1}{3}$                       (b)  $\frac{2}{3}$                       (c)  $\frac{1}{4}$                       (d)  $\frac{3}{4}$

Q60. The Hamiltonian of a one-dimensional Ising model of  $N$  spins ( $N$  large) is

$$H = -J \sum_{i=1}^N \sigma_i \sigma_{i+1}$$


where the spin  $\sigma_i = \pm 1$  and  $J$  is a positive constant. At inverse temperature  $\beta = \frac{1}{k_B T}$ , the correlation function between the nearest neighbor spins ( $\sigma_i \sigma_{i+1}$ ) is

- (a)  $\frac{e^{-\beta J}}{e^{\beta J} + e^{-\beta J}}$  (b)  $e^{-2\beta J}$   
 (c)  $\tan h(\beta J)$  (d)  $\cot h(\beta J)$

Q61. At low temperatures, in the Debye approximation, the contribution of the phonons to the heat capacity of a two dimensional solid is proportional to

- (a)  $T^2$  (b)  $T^3$  (c)  $T^{1/2}$  (d)  $T^{3/2}$

Q62. A particle hops on a one-dimensional lattice with lattice spacing  $a$ . The probability of the particle to hop to the neighboring site to its right is  $p$ , while the corresponding probability to hop to the left is  $q = 1 - p$ . The root-mean squared deviation  $\Delta x = \sqrt{\langle x \rangle^2 - \langle x \rangle^2}$  in displacement after  $N$  steps, is




- (a)  $a\sqrt{Npq}$  (b)  $aN\sqrt{pq}$  (c)  $2a\sqrt{Npq}$  (d)  $a\sqrt{N}$

Q63. The energy levels accessible to a molecule have energies  $E_1 = 0, E_2 = \Delta$  and  $E_3 = 2\Delta$  (where  $\Delta$  is a constant). A gas of these molecules is in thermal equilibrium at temperature  $T$ . The specific heat at constant volume in the high temperature limit ( $k_B T \gg \Delta$ ) varies with temperature as





- (a)  $\frac{1}{T^{3/2}}$  (b)  $\frac{1}{T^3}$  (c)  $\frac{1}{T}$  (d)  $\frac{1}{T^2}$

Q64. The input  $V_i$  to the following circuit is a square wave as shown in the following figure.







which of the waveforms best describes the output?





Q65. The amplitude of a carrier signal of frequency  $f_0$  is sinusoidally modulated at a frequency  $f' \ll f_0$ . Which of the following graphs best describes its power spectrum?





Q66. The standard deviation of the following set of data

$\{10.0, 10.0, 9.9, 9.9, 9.8, 9.9, 9.9, 9.9, 9.9, 9.8, 9.9\}$  is nearest to

- (a) 0.10                      (b) 0.07                      (c) 0.01                      (d) 0.04

Q67. The diatomic molecule HF has an absorption line in the rotational band at  $40 \text{ cm}^{-1}$  for the isotope  $^{18}\text{F}$ . The corresponding line for the isotope  $^{19}\text{F}$  will be shifted by approximately

- (a)  $0.05 \text{ cm}^{-1}$               (b)  $0.11 \text{ cm}^{-1}$               (c)  $0.33 \text{ cm}^{-1}$               (d)  $0.01 \text{ cm}^{-1}$

Q68. The excited state ( $n = 4, l = 2$ ) of an electron in an atom may decay to one or more of the lower energy levels shown in the diagram below.

$$n = 4 \quad \overline{\quad} \\ l = 2$$

$$n = 3 \quad \overline{\quad} \quad \overline{\quad} \quad \overline{\quad} \\ l = 0 \quad l = 1 \quad l = 2$$

$$n = 2 \quad \overline{\quad} \\ l = 1$$

Of the total emitted light, a fraction  $\frac{1}{4}$  comes from the decay to the state ( $n = 2, l = 1$ ). Based on selection rules, the fractional intensity of the emission line due to the decay to the state ( $n = 3, l = 1$ )


- (a)  $\frac{3}{4}$                       (b)  $\frac{1}{2}$                       (c)  $\frac{1}{4}$                       (d) 0

Q69. The volume of an optical cavity is  $1 \text{ cm}^3$ . The number of modes it can support within a bandwidth of  $0.1 \text{ nm}$ , centered at  $\lambda = 500 \text{ nm}$ , is of the order of

- (a)  $10^3$                       (b)  $10^5$                       (c)  $10^{10}$                       (d)  $10^7$

Q70. Barium Titanate ( $\text{BaTiO}_3$ ) crystal has a cubic perovskite structure, where the  $\text{Ba}^{2+}$  ions are at the vertices of a unit cube, the  $\text{O}^{2-}$  ions are at the centres of the faces while the  $\text{Ti}^{2+}$  is at the centre. The number of optical phonon modes of the crystal is

- (a) 12    (b) 15  
(c) 5    (d) 18



Q71. The dispersion relation of optical phonons in a cubic crystal is given

by  $\omega(k) = \omega_0 - ak^2$  where  $\omega_0$  and  $a$  are positive constants. The contribution to the density of states due to these phonons with frequencies just below  $\omega_0$  is proportional to

- (a)  $(\omega_0 - \omega)^{1/2}$                       (b)  $(\omega_0 - \omega)^{3/2}$                       (c)  $(\omega_0 - \omega)^2$                       (d)  $(\omega_0 - \omega)$

