

GATE PHYSICS 2023 SOLUTION

General Aptitude

Q1. – Q5. Carry ONE mark each.

Q.1 "You are delaying the completion of the task. Send contributions at the earliest."

- (A) you are (B) your (C) you're (D) yore

Ans. : (B)

Q.2 References :..... : Guidelines : Implement

(By word meaning)

- (A) Sight (B) Site (C) Cite (D) Plagiarise

Ans. : (C)


Q.3 In the given figure, PQRS is a parallelogram with $PS = 7$ cm, $PT = 4$ cm and $PV = 5$ cm. What is the length of RS in cm ? (The diagram is representative.)

(A) $\frac{20}{7}$

(B) $\frac{28}{5}$

(C) $\frac{9}{2}$

(D) $\frac{35}{4}$



Ans. : (B)

Q.4 In 2022, June Huh was awarded the Fields medal, which is the highest prize in Mathematics.

When he was younger, he was also a poet. He did not win any medals in the International Mathematics Olympiads. He dropped out of college.

Based only on the above information, which one of the following statements can be logically inferred with certainty?

(A) Every Fields medalist has won a medal in an International Mathematics Olympiad.

(B) Everyone who has dropped out of college has won the Fields medal.


(C) All Fields medalists are part-time poets.

(D) Some Fields medalists have dropped out of college.

Ans. : (D)

Q.5 A line of symmetry is defined as a line that divides a figure into two parts in a way such that each part is a mirror image of the other part about that line.

The given figure consists of 16 unit squares arranged as shown. In addition to the three black squares, what is the minimum number of squares that must be coloured black, such that both PQ and MN form lines of symmetry? (The figure is representative)



- (A) 3 (B) 4 (C) 5 (D) 6

Ans. : (C)

Q.6 - Q.10 Carry TWO marks Each

Q.6 Human beings are one among many creatures that inhabit an imagined world. In this imagined world, some creatures are cruel. If in this imagined world, it is given that the statement "Some human beings are not cruel creatures" is FALSE, then which of the following set of statement(s) can be logically inferred with certainty?

- (i) All human beings are cruel creatures.
 - (ii) Some human beings are cruel creatures.
 - (iii) Some creatures that are cruel are human beings.
 - (iv) No human beings are cruel creatures.
- (A) only (i) (B) only (iii) and (iv)
 (C) only (i) and (ii) (D) (i), (ii) and (iii)

Ans. : (D)

Q.7 To construct a wall, sand and cement are mixed in the ratio of 3: 1. The cost of sand and that of cement are in the ratio of 1: 2.

If the total cost of sand and cement to construct the wall is 1000 rupees, then what is the cost (in rupees) of cement used?

- (A) 400 (B) 600 (C) 800 (D) 200

Ans. : (A)

Q.8 The World Bank has declared that it does not plan to offer new financing to Sri Lanka, which is battling its worst economic crisis in decades, until the country has an adequate macroeconomic policy framework in place. In a statement, the World Bank said Sri Lanka needed to adopt structural reforms that focus on economic stabilisation and tackle the root causes of its crisis. The latter has starved it of foreign exchange and led to shortages of food, fuel, and medicines. The bank is repurposing resources under existing loans to help alleviate shortages of essential items such as medicine, cooking gas, fertiliser, meals for children, and cash for vulnerable households. Based only on the above passage, which one of the following statements can be inferred with certainty?

(A) According to the World Bank, the root cause of Sri Lanka's economic crisis is that it does not have enough foreign exchange.

(B) The World Bank has stated that it will advise the Sri Lankan government about how to tackle the root causes of its economic crisis.

(C) According to the World Bank, Sri Lanka does not yet have an adequate macroeconomic policy framework.

(D) The World Bank has stated that it will provide Sri Lanka with additional funds for essentials such as food, fuel, and medicines.

Ans. : (C)

Q.9 The coefficient of x^4 in the polynomial $(x - 1)^3(x - 2)^3$ is equal to.....

(A) 33

(B) -3

(C) 30

(D) 21

Ans. : (A)

Q.10 Which one of the following shapes can be used to tile (completely cover by repeating) a flat plane, extending to infinity in all directions, without leaving any empty spaces in between them? The copies of the shape used to tile are identical and are not allowed to overlap.

(A) circle

(B) regular octagon






(C) regular pentagon

(D) rhombus

Ans. : (D)

Q.11 - Q.35 Carry ONE mark Each

Q.11 Which one of the following entropy (S) - temperature (T) diagrams CORRECTLY represents the Carnot cycle $abcda$ shown in the $P - V$ diagram?



Topic: Thermodynamics and Statistical Mechanics

Sub-Topic: Entropy

Ans. (A)

Solution: a to b should be a constant temperature so B and D is in correct C is incorrect because b to c should be adiabatic a constant entropy curve.

Q.12 Which one of the following is a dimensionless constant?

- | | |
|--------------------------------|--------------------------------|
| (A) Permittivity of free space | (B) Permeability of free space |
| (C) Bohr magneton | (D) Fine structure constant |

Topic: Atomic, Molecular and Laser Physics

Sub-Topic: Fine structure

Ans. : (D)

Solution: Fine structure constant

$$\alpha = \frac{\text{Speed of electron in first Bohr Orbit}}{\text{Speed of electron in vacuum}} = \frac{LT^{-1}}{LT^{-1}} = \text{Dimensionless}$$

Q.13 Choose the most appropriate matching of the items in Column 1 with those in Column 2

| Column 1 | Column 2 |
|-------------------|------------------------------------------|
| (i) PIN diode | P. Voltage regulation |
| (ii) Tunnel diode | Q. Radio frequency and microwave devices |
| (iii) Zener diode | R. Optoelectronic detection |
| (iv) Photo diode | S. Oscillator |

(A) (i) - Q; (ii) - S; (iii) - P; (iv) - R

(B) (i) - R; (ii) - Q; (iii) - P; (iv) - S

(C) (i) - R; (ii) - S; (iii) - P; (iv) - Q

(D) (i) - P; (ii) - Q; (iii) - R; (iv) - S

Topic: Electronics

Sub-Topic: Diode

Ans. : (A)

Q.14 The atomic number of an atom is 6 . What is the spectroscopic notation of its ground state, according to Hund's rules?

(A) 3P_0

(B) 3P_1

(C) 3D_3

(D) 3S_1

Topic: Atomic, Molecular and Laser Physics

Sub-Topic: L-S Coupling

Ans. : (A)

Solution: Electronic configuration of atomic number 6 (Carbon)

$$1s^2 2s^2 2p^2$$

Optical electrons are equivalent with less than half filled, these are $2p^2$

$$2p^2 \Rightarrow l_1 = 1, l_2 = 1$$

$$l_1 = 1, l_2 = 1$$

$$s_1 = \frac{1}{2}, s_2 = \frac{1}{2}$$

$$\text{Maximize } L = |m_{l1}| + |m_{l2}| = 1 + 0 = 1, \text{ Maximize } S = |m_{s1}| + |m_{s2}| = 1$$

$$\text{Less than half filled } J = L - S = 0$$

$$\text{G.S } ^3P_0$$

Ans. : (B)

Solution: For rotational Matrix $\det A = 1$.

Q.18 A compound consists of three ions X, Y and Z. The Z ions are arranged in an FCC arrangement. The X ions occupy $\frac{1}{6}$ of the tetrahedral voids and the Y ions occupy $\frac{1}{3}$ of the octahedral voids. Which one of the following is the CORRECT chemical formula of the compound?

- (A) XY_2Z_4 (B) XYZ_3 (C) XYZ_2 (D) XYZ_4

Topic: Condensed Matter Physics

Sub-Topic: Crystal Structure

Ans. : (B)

Solution: Effective lattice point in FCC unit cell =4,

No of Z ions in the unit cell =4

No. of tetrahedral voids =8

No of X ions in the unit cell = $\frac{8}{6} = \frac{4}{3}$

No. of octahedral voids =4

No of Y ions in the unit cell = $\frac{4}{3}$





Chemical compound ratio of ions, $\frac{4}{3} : \frac{4}{3} : 4 = 1 : 1 : 3$,

Formula of Chemical compound = XYZ_3

Topic: Condensed Matter Physics

Sub-Topic: Crystal Structure

Q.19 For a non-magnetic metal, which one of the following graphs best represents the behavior of $\frac{C}{T}$ vs. T^2 , where C is the heat capacity and T is the temperature?



Topic: Condensed Matter Physics

Sub-Topic: Heat Capacity

Ans. : (B)

Solution: For non -magnetic metal $C = AT + BT^3 \Rightarrow \frac{C}{T} = A + BT^2$

Ans. : (B)

Solution: For type I Superconductor $\chi = \frac{M}{H} = -1$

For diamagnetic material χ is small & negative

For paramagnetic material χ is small & positive


Q.22 Graphene is a two-dimensional material, in which carbon atoms are arranged in a honeycomb lattice with lattice constant a . As shown in the figure, \vec{a}_1 and \vec{a}_2 are two lattice vectors. Which one of the following is the area of the first Brillouin zone for this lattice?

(A) $\frac{8\pi^2}{3\sqrt{3}a^2}$

(B) $\frac{4\pi^2}{3\sqrt{3}a^2}$

(C) $\frac{8\pi^2}{\sqrt{3}a^2}$

(D) $\frac{4\pi^2}{\sqrt{3}a^2}$



Topic: Solid state Physics

Sub-Topic: Crystal structure


Ans. : (A)

Solution: Area of the cell $A = 6 \times \frac{\sqrt{3}}{4} a^2 = \frac{3\sqrt{3}}{2} a^2$

Area of the 1st Brillouin zone

$$A^* = \frac{(2\pi)^2}{A} = \frac{4\pi^2}{\frac{3\sqrt{3}}{2} a^2} = \frac{8\pi^2}{3\sqrt{3}a^2}$$

Q.23 ^{60}Co nucleus emits a β -particle and is converted to $^{60}\text{Ni}^*$ with $J^P = 4^+$, which in turn decays to the ^{60}Ni ground state with $J^P = 0^+$ by emitting two photons in succession, as shown in the figure. Which one of the following statements is CORRECT?



- (A) $4^+ \rightarrow 2^+$ is an electric octupole transition
- (B) $4^+ \rightarrow 2^+$ is a magnetic quadrupole transition
- (C) $2^+ \rightarrow 0^+$ is an electric quadrupole transition
- (D) $2^+ \rightarrow 0^+$ is a magnetic quadrupole transition

Topic: Nuclear and Particle Physics


Sub-Topic: Radioactivity

Ans. : (C)

Solution: $4^+ \rightarrow 2^+ \Rightarrow |\vec{J}_i - \vec{J}_f| \leq L \leq |\vec{J}_i + \vec{J}_f|, \Delta\pi = \text{No change} \ \& \ L = 2,3,4,5,6$

So, elects $4^+ \rightarrow 2^+$ transition will be of electric quadrupole (E2) type other magnetic multipole transitions are possible here. Thus, contribution/Intensity of E2 will be less. Here, $2^+ \rightarrow 0^+$ ($\Delta\pi =$ No change, $L = 2$). So, $2^+ \rightarrow 0^+$ transition will only be of electric quadrupole (E2) type. Therefore, it will be highly intense.

Q.24 Which one of the following options is CORRECT for the given logic circuit?



- (A) $P = 1, Q = 1; X = 0$ (B) $P = 1, Q = 0; X = 1$
 (C) $P = 0, Q = 1; X = 0$ (D) $P = 0, Q = 0; X = 1$

Topic: Electronics

Sub-Topic: Logic gate

Ans. : (D)

Solution: By using binary algebra, $X = \bar{P} + PQ = \bar{P} + \bar{P}Q = \bar{P} + Q$

$$P = 0, Q = 0 \Rightarrow X = \bar{0} + 0 = 1 + 0 = 1$$

Q.25 An atom with non-zero magnetic moment has an angular momentum of magnitude $\sqrt{12}\hbar$. When a beam of such atoms is passed through a SternGerlach apparatus, how many beams does it split into?

- (A) 3 (B) 7 (C) 9 (D) 25

Topic: Quantum Mechanics

Sub-Topic: Spin

Ans. : (B)

Solution: $s(s+1) = 12 \Rightarrow s = 3$ number of states is $2s+1 = 2.3+1 = 7$

Q.26 A 4×4 matrix M has the property $M^\dagger = -M$ and $M^4 = \mathbf{1}$, where $\mathbf{1}$ is the 4×4 identity matrix. Which one of the following is the CORRECT set of eigenvalues of the matrix M ?

- (A) $(1, 1, -1, -1)$ (B) $(i, i, -i, -i)$ (C) $(i, i, i, -i)$ (D) $(1, 1, -i, -i)$

Topic: Mathematical Physics

Sub-Topic: Matrix

Ans. : (B)

Solution: It is a skew Hermitian matrix. So, Eigen values are imaginary or zero. Moreover, Trace should be zero. Also $\sum_i \lambda_i^4 = 4$

Ans. : (A) and (C)

Solution: A is correct because $Q = \frac{2}{5}Z(b^2 - a^2) \neq 0, \because b \neq a,$

Due to non-spherical and even parity of deuteron. The ground state is mixing of 3S_1 & 3D_1

Q.30 The Geiger-Muller counter is a device to detect α, β and γ radiations. It is a cylindrical tube filled with monatomic gases like argon, and polyatomic gases such as ethyl alcohol. The inner electrode is along the axis of the cylindrical tube and the outer electrode is the tube. Which of the following statements is(are) CORRECT?

- (A) Argon is used so that ambient light coming from the surroundings do not produce any signal in the detector
- (B) Ethyl alcohol is used as a quenching gas
- (C) The electric field strength decreases from the axis to the edge of the tube and the direction of the field is radially outward
- (D) The electric field increases from the axis to the edge of the tube and the field direction is radially inward

Topic: Nuclear and Particle Physics
Sub-Topic: Nuclear Detector

Ans. : (A), (B) and (C)

Solution: The basic construction and working principle of Geiger-Muller demonstrated that

Argon is used so that ambient light coming from the surroundings do not produce any signal in the detector. Ethyl alcohol is used as quenching vapour. Electric field at a distance from r of the center

$$E = \frac{V}{r \ln \frac{b}{a}}$$

Q.31 Consider an isolated magnetized sphere of radius R with a uniform magnetization \vec{M} along the positive z direction, with the north and south poles of the sphere lying on the Z axis. It is given that the magnetic field inside the sphere is $\vec{B} = \frac{2\mu_0}{3}\vec{M}$, where μ_0 is the permeability of vacuum. Which of the following statements is(are) CORRECT?

- (A) The bound volume current density is zero
- (B) The bound surface current density has maximum magnitude at the equator, where this magnitude equals $|\vec{M}|$
- (C) The auxiliary field $\vec{H} = -\frac{2}{3}\vec{M}$
- (D) Far from the sphere, the magnetic field is due to a dipole of moment \vec{m} , (where $\frac{\vec{m}}{4\pi R^3} = \frac{B}{2\mu_0}\hat{Z}$)

Topic: EMT
Sub-Topic: Magnetostatic

Ans. : (A), (B) and (D)

Solution: (A) $J_b = \nabla \times M = 0$ (Since M is uniform)

(B) $\vec{K}_b = \vec{M} \times \hat{n} = M \sin \theta \hat{\phi}$, So, At the equator $\theta = 0$ and $|\vec{K}_b| = M$

(C) $\vec{H} = \frac{\vec{B}}{\mu_0} - \vec{M} = \frac{2\mu_0}{3\mu_0} M - M = -\frac{1}{3} \vec{M}$

(D) $B = \frac{\mu_0 m}{4\pi r^3} [2 \cos \theta \hat{r} + \sin \theta \hat{\theta}]$, To relate dipole moment, take theta zero then $B = \frac{\mu_0 m}{4\pi r^3}$,

$$\frac{\vec{m}}{4\pi R^3} = \frac{B}{2\mu_0} \hat{z}.$$

Far from the sphere field will be non-uniform and will be due to dipole moment \vec{m} .

Q.32 Which of the following options represent(s) linearly independent pair(s) of functions of a real variable x ?

- (A) e^{ix} and e^{-ix} (B) x and e^x (C) 2^x and 2^{-3+x} (D) e^{ix} and $\sin x$

Topic: Mathematical Physics

Sub-Topic: Differential Equation

Ans. : (A), (B) and (D)

Solution: Let us check the Wronskian of the solutions

(A) $W = \begin{vmatrix} e^{ix} & e^{-ix} \\ ie^{ix} & -ie^{-ix} \end{vmatrix} = -i - i = -2i \neq 0$ Linearly Independent

(B) $W = \begin{vmatrix} x & e^x \\ 1 & e^x \end{vmatrix} = xe^x - e^x \neq 0$ Linearly Independent

(C) $W = \begin{vmatrix} 2^x & 2^{-3+x} = 2^{-3} \cdot 2^x \\ 2^x \ln 2 & 2^{-3} \cdot 2^x \ln 2 \end{vmatrix} = 2^{-3} 2^x \cdot 2^x \ln 2 - 2^{-3} 2^x \cdot 2^x \ln 2 = 0$ Linearly dependent

(D) $W = \begin{vmatrix} 3^{ix} & \sin x \\ ie^{ix} & \cos x \end{vmatrix} = e^{ix} \cos x - ie^{ix} \sin x \neq 0$ Linearly Independent

Q.33 In the vector model of angular momentum applied to atoms, what is the minimum angle in degrees (in integer) made by the orbital angular momentum vector and the positive Z axis for a $2p$ electron?

Topic: Atomic, Molecular and Laser Physics

Sub-Topic: Vector Model

Ans. : 45°

Solution: Given for $2p$, $\cos \theta = \frac{m_l}{\sqrt{l(l+1)}}$, $2p = 1, m_l = 1, 0, -1$, $\cos \theta = \frac{1}{\sqrt{2}}, 0, -\frac{1}{\sqrt{2}}$, $\theta = 45, 90, 135$

Q.34 For a transistor amplifier, the frequency response is such that the mid band voltage gain is 200. The cutoff frequencies are 20 Hz and 20kHz. What is the ratio (rounded off to two decimal places) of the voltage gain at 10 Hz to that at 100 kHz ?

Topic: Electronics

Sub-Topic: Amplifier

Ans. : 2.28

Solution: For 10 Hz

$$A1(f) = \frac{A}{\sqrt{1 + \left(\frac{f_L}{f}\right)^2}} = \frac{200}{\sqrt{1 + \left(\frac{20}{10}\right)^2}} = \frac{200}{\sqrt{5}} = 89.44$$

For 100 Hz

$$A2(f) = \frac{A}{\sqrt{1 + (f/f_H)^2}} = \frac{200}{\sqrt{1 + \left(\frac{100}{20}\right)^2}} = 39.22$$

The ratio of gain will be $= \frac{89.44}{39.22} = 2.28$

Q.35 An electric field as a function of radial coordinate r has the form $\vec{E} = \alpha \frac{e^{-r^2}}{r} \hat{r}$, where α is a constant. Assume that dimensions are appropriately taken care of. The electric flux through a sphere of radius $\sqrt{2}$, centered at the origin, is Φ . What is the value of $\frac{\Phi}{2\pi\alpha}$ (rounded off to two decimal places)?

Topic: EMT

Sub-Topic: Electrostatic

Ans. : 0.38

Solution: $\phi = \int \vec{E} \cdot d\vec{a} = \alpha \frac{e^{-R^2}}{R} \times 4\pi R^2, \phi = 4\pi R\alpha e^{-R^2} \Rightarrow \frac{\phi}{2\pi\alpha} = \frac{4\pi R\alpha e^{-R^2}}{2\pi\alpha} = 2R e^{-R^2} = 2\sqrt{2} e^{-2} = 0.38$

Q.36 - Q.65 Carry TWO marks Each

Q.36 It is given that the electronic ground state of a diatomic molecule X_2 has even parity and the nuclear spin of X is 0. Which one of the following is the CORRECT statement with regard to the rotational Raman spectrum (J is the rotational quantum number) of this molecule?

- (A) Lines of all J values are present
- (B) Lines have alternating intensity in the ratio of 3:1
- (C) Lines of only even J values are present
- (D) Lines of only odd J values are present






Atomic, Molecular and Laser Physics

Sub-Topic: Raman Effect

Ans.: (C)

Solution: For given molecule, the nuclear wavefunction corresponding to odd J is anti-symmetric and it is symmetric for even J . Also, the degeneracy of anti-symmetric (Odd J) $(2I + 1)I = 0$. Also, the degeneracy of anti-symmetric $(2I + 1)(I + 1) = 1$. Thus, odd J states will be absent. Only, even J states will be present. Thus, lines for even J will be present.

Q.37 An input voltage in the form of a square wave of frequency 1kHz is given to a circuit, which results in the output shown schematically below. Which one of the following options is the CORRECT representation of the circuit?



Topic: Electronics

Sub-Topic: RC Circuit

Ans. : (A)

Solution: Input signal will be differentiated.

For differentiated output

$$RC \geq 5T$$

$$RC = 5 \times 10^3 \times 1 \times 10^{-6} = 5\text{msec}$$

$$T = \frac{1}{f} = \frac{1}{10^3} = 1\text{msec}$$

Q38. A simple harmonic oscillator with an angular frequency ω is in thermal equilibrium with a reservoir at absolute temperature T , with $\omega = \frac{2k_B T}{\hbar}$. Which one of the following is the partition function of the system?

Topic: Thermodynamics and Statistical Mechanics

Sub-Topic: Partition Function

Ans. : (A)

$$\text{Solution: } z_1 = \frac{e^{-\frac{1}{2}\beta\hbar\omega}}{1 - e^{-\beta\hbar\omega}} = \frac{e^{-\frac{1}{2} \frac{\hbar}{2k_B T} \left(\frac{2k_B T}{\hbar} \right)}}{1 - e^{-2}} = \frac{e^{-1}}{1 - e^{-2}} = \frac{1}{e - e^{-1}} = \frac{e}{e^2 - 1}$$

Q.39 Which one of the following options is the most appropriate match between the items given in Column 1 and Column 2?

| Column 1 | Column 2 |
|-----------------------|---------------------------------------------------|
| (i) Visible light | P. Transition between core energy levels of atoms |
| (ii) X-rays | Q. Transition between nuclear energy levels |
| (iii) Gamma rays | R. Pair production |
| (iv) Thermal neutrons | S. Crystal structure determination |
| | T. Photoelectric effect |

- (A) (i) - T; (ii) - P, S, T; (iii) - Q, R; (iv) - S
- (B) (i) - P, T; (ii) - S; (iii) - R, S; (iv) - S, T
- (C) (i) - T; (ii) - R, S; (iii) - Q, R; (iv) - S
- (D) (i) - S, T; (ii) - P, S; (iii) - R, T; (iv) - S


Topic: Atomic, Molecular and Laser Physics

Sub-Topic: Molecular Physics

Ans. : (A)

Solution: Visible light can only do Photoelectric effect because it has very less energy which cannot do the other given process. X-Ray has enough energy which can make transition from inner core electron, Crystal structure, Photoelectric effect. Gamma ray can do transition between electronic level and pair production, these are well known from basic concept of radioactivity. Thermal neutron can do crystal structure determination.

- Q.40 A rod PQ of proper length L lies along the X -axis and moves towards the positive x direction with speed $v = \frac{3c}{5}$ with respect to the ground (see figure), where c is the speed of light in vacuum. An observer on the ground measures the positions of P and Q at different times t_P and t_Q respectively in the ground frame, and finds the difference between them to be $\frac{9L}{10}$. What is the value of $t_Q - t_P$?



- (A) $\frac{L}{3c}$ (B) $\frac{L}{5c}$ (C) $\frac{L}{6c}$ (D) $\frac{2L}{3c}$

Topic: Classical Mechanics

Sub-Topic: STR

Ans. : (C)

Solution: $\Delta t = t_Q - t_P = \frac{\Delta t'}{\sqrt{1 - \frac{v^2}{c^2}}} + \frac{\Delta x' \frac{v}{c^2}}{\sqrt{1 - \frac{v^2}{c^2}}}$ $\Delta x' = L, v = \frac{3c}{5}$

It is also given

$$\Delta x = x_Q - x_P = \frac{9L}{10} = \frac{\Delta x'}{\sqrt{1 - \frac{v^2}{c^2}}} + \frac{\Delta t' v}{\sqrt{1 - \frac{v^2}{c^2}}} \Rightarrow \frac{\Delta t'}{\sqrt{1 - \frac{v^2}{c^2}}} = -\frac{7}{12} \frac{L}{c}$$

$$\Delta t = t_Q - t_P = -\frac{7}{12} \frac{L}{c} + \frac{3}{5} \frac{5L}{4} \Rightarrow \frac{L}{6c}$$

- Q.41 A symmetric top has principal moments of inertia $I_1 = I_2 = \frac{2\alpha}{3}, I_3 = 2\alpha$ about a set of principal axes 1,2,3 respectively, passing through its center of mass, where α is a positive constant. There is no force acting on the body and the angular speed of the body about the 3-axis is $\omega_3 = \frac{1}{8}$ rad/s. With what angular frequency in rad/s does the angular velocity vector $\vec{\omega}_1$ precess about the 3-axis?
- (A) 2 (B) 3 (C) 5 (D) 7

Topic: Classical Mechanics

Sub-Topic: Moment of inertia

Ans.: MTA

Q.42 A particle of mass m is free to move on a frictionless horizontal two dimensional (r, θ) plane, and is acted upon by a force $\vec{F} = -\frac{k}{2r^3}\hat{r}$ with k being a positive constant. If p_r and p_θ are the generalised momenta corresponding to r and θ respectively, then what is the value of $\frac{dp_r}{dt}$?

- (A) $\frac{p_\theta^2 - 2mk}{2mr^3}$ (B) $\frac{2p_\theta^2 - mk}{mr^3}$ (C) $\frac{p_\theta^2 - 2mk}{mr^3}$ (D) $\frac{2p_\theta^2 - mk}{2mr^3}$

Topic: Classical Mechanics

Sub-Topic: Hamiltonian

Ans. : (D)

Solution: Force $F = -\frac{\partial V}{\partial r}\hat{r} = -\frac{k}{2r^3}\hat{r}$

If motion is in two-dimensional plane, then p_r and p_θ are generalized momentum for generalized coordinates conjugate to r and θ . If $V(r)$ is potential then Hamiltonian in polar coordinate is given by $H = \frac{p_r^2}{2m} + \frac{p_\theta^2}{2mr^2} + V(r)$. The Hamiltonian equation of motion is given by

$$\frac{\partial H}{\partial r} = -\dot{p}_r \Rightarrow -\frac{p_\theta^2}{mr^3} + \frac{\partial V}{\partial r} \Rightarrow -\frac{p_\theta^2}{mr^3} + \frac{k}{2r^3} = -\frac{dp_r}{dt} \Rightarrow \frac{dp_r}{dt} = \frac{2p_\theta^2 - mk}{2mr^3}$$

Q.43 Consider two real functions

$$U(x, y) = xy(x^2 - y^2)'$$

$$V(x, y) = ax^4 + by^4 + cx^2y^2 + k,$$

where k is a real constant and a, b, c are real coefficients. If $U(x, y) + iV(x, y)$ is analytic, then what is the value of $a \times b \times c$

- (A) $\frac{1}{8}$ (B) $\frac{3}{28}$ (C) $\frac{5}{36}$ (D) $\frac{3}{32}$

Topic: Mathematical Physics

Sub-Topic: Function of two variables

Ans. : (D)

Solution: $U_x = V_y, 3x^2y - y^3 = 4by^3 + 2cx^2y, 4b = -1, 2c = 3, U_y = -V_x$

$$x^3 - 3xy^2 = -(4ax^3 + 2cx^2y), -4a = 1, a = -\frac{1}{4}, b = -\frac{1}{4}, c = \frac{3}{2}$$

$$\Rightarrow -\frac{1}{4} \times -\frac{1}{4} \times \frac{3}{2} = \frac{3}{32}$$

- Q.44 Young's double slit experiment is performed using a beam of C_{60} (fullerene) molecules, each molecule being made up of 60 carbon atoms. When the slit separation is 50 nm, fringes are formed on a screen kept at a distance of 1 m from the slits. Now, the experiment is repeated with C_{70} molecules with a slit separation of 92.5 nm. The kinetic energies of both the beams are the same. The position of the 4th bright fringe for C_{60} will correspond to the n^{th} bright fringe for C_{70} . What is the value of n (rounded off to the nearest integer)?
- (A) 5 (B) 6 (C) 7 (D) 8

Topic: Optics

Sub-Topic: Interference

Ans. : (D)

Solution: YDSE with C_{60} beam

$$\lambda_1 = \frac{h}{\sqrt{2mk}} = \frac{h}{\sqrt{2 \times (60 \times 12\text{amu})k}}$$

YDSE with C_{70} beam

$$\lambda_2 = \frac{h}{\sqrt{2 \times (70 \times 12\text{amu})k}}$$

$$\beta_2 = \frac{D\lambda_2}{d_2} = \frac{1 \text{ m}}{92.5 \text{ nm}} \frac{h}{\sqrt{2 \times (70 \times 12\text{amu})k}}$$

Given $n\beta_2 = 4\beta_1$

$$\frac{n}{92.5\sqrt{70}} = \frac{4}{50\sqrt{60}} \Rightarrow n = 8$$

- Q.45 A neutron beam with a wave vector \vec{k} and an energy 20.4meV diffracts from a crystal with an outgoing wave vector \vec{k}' . One of the diffraction peaks is observed for the reciprocal lattice vector \vec{G} of magnitude 3.14\AA^{-1} . What is the diffraction angle in degrees (rounded off to the nearest integer) that \vec{k} makes with the plane? (Use mass of neutron = $1.67 \times 10^{-27}\text{Kg}$)
- (A) 15 (B) 30 (C) 45 (D) 60

Topic: Condensed Matter Physics

Sub-Topic: Bragg's Law

Ans.: (B)

Solution: Energy of neutron beam = 20.4meV

Corresponding de Broglie wavelength

$$\lambda = \frac{h}{p} = \frac{h}{\sqrt{2mE}} = \frac{6.62 \times 10^{-34}}{\sqrt{2 \times 1.67 \times 10^{-27} \times 20.4 \times 10^{-3} \times 1.6 \times 10^{-19}}} = 2\text{\AA}$$

Diffraction condition

$$2d \sin \theta = \lambda \quad \& \quad = \frac{2\pi}{\|G\|}$$

$$\sin \theta = \lambda \frac{\|G\|}{4\pi} = \frac{2 \times \text{\AA} \times 3.14 \text{\AA}^{-1}}{4\pi} = \frac{1}{2}$$

$$\theta = 30^\circ$$

Q.46 In the first Brillouin zone of a rectangular lattice (lattice constants $a = 6\text{\AA}$ and $b = 4\text{\AA}$), three incoming phonons with the same wave vector $\langle 1.2\text{\AA}^{-1}, 0.6\text{\AA}^{-1} \rangle$ interact to give one phonon.

Which one of the following is the CORRECT wave vector of the resulting phonon?

- (A) $\langle 2.56\text{\AA}^{-1}, 0.23\text{\AA}^{-1} \rangle$ (B) $\langle 3.60\text{\AA}^{-1}, 1.80\text{\AA}^{-1} \rangle$
 (C) $\langle 0.48\text{\AA}^{-1}, 0.23\text{\AA}^{-1} \rangle$ (D) $\langle 3.60\text{\AA}^{-1}, -0.80\text{\AA}^{-1} \rangle$

Topic: Solid state Physics

Sub-Topic: Brillouin zone

Ans. : (A)

Solution: $\vec{k}_1 + \vec{k}_2 + \vec{k}_3 = \vec{k} + \vec{G} \Rightarrow \vec{k} = \vec{k}_1 + \vec{k}_2 + \vec{k}_3 - \vec{G}$

$$\therefore k_x = k_{1x} + k_{2x} + k_{3x} - G_x \quad \text{and} \quad k_y = k_{1y} + k_{2y} + k_{3y} - G_y$$

$$\text{where } G_x = \frac{2\pi}{a} = \frac{2\pi}{6} = \frac{\pi}{3} A^{0-1} = 1.047 A^{0-1} \quad \& \quad G_y = \frac{2\pi}{b} - \frac{2\pi}{4} = \frac{\pi}{2} A^{0-1} = 1.57 A^{0-1}$$

Thus,

$$k_x = 1.2 + 1.2 + 1.2 - 1.047 = 2.56 A^{0-1} \quad \& \quad k_y = 0.6 + 0.6 + 0.6 - 1.57 = 0.23 A^{0-1}$$

Q.47 For a covalently bonded solid consisting of ions of mass m , the binding potential can be assumed to be given by

$$U(r) = -\epsilon \left(\frac{r}{r_0} \right) e^{-\frac{r}{r_0}}$$

where ϵ and r_0 are positive constants. What is the Einstein frequency of the solid in Hz ?

- (A) $\frac{1}{2\pi} \sqrt{\frac{\epsilon\epsilon}{mr_0^2}}$ (B) $\frac{1}{2\pi} \sqrt{\frac{\epsilon}{\text{mer}_0^2}}$
 (C) $\frac{1}{2\pi} \sqrt{\frac{2\epsilon}{\text{mer}_0^2}}$ (D) $\frac{1}{2\pi} \sqrt{\frac{\epsilon\epsilon}{2mr_0^2}}$

Topic: Condensed Matter Physics

Sub-Topic: Einstein Theory

Ans. : (B)

Solution: From the concept of stability analysis





$$\frac{du}{dr} = 0 \Rightarrow \left[e^{-\frac{r}{r_0}} - \frac{r}{r_0} e^{-\frac{r}{r_0}} \right] = 0$$

$$\Rightarrow e^{-\frac{r}{r_0}} \left[1 - \frac{r}{r_0} \right] = 0 \Rightarrow r = r_0$$

$$\text{Now, } \frac{d^2u}{dr^2} = -\frac{\epsilon}{r_0} e^{-\frac{r}{r_0}} \left[-\frac{r_0}{r_0} + \frac{r}{r_0^2} \right] = \frac{\epsilon}{er_0^2} \text{ (At, } r = r_0 \text{)}$$

$$f = \frac{1}{2\pi} \sqrt{\frac{d^2u}{dr^2} \bigg|_{r=r_0}} = \frac{1}{2\pi} \sqrt{\frac{\epsilon}{mer_0^2}}$$

Q.48 In a hadronic interaction, π^0 's are produced with different momenta, and they immediately decay into two photons with an opening angle θ between them. Assuming that all these decays occur in one plane, which one of the following figures depicts the behaviour of θ as a function of the π^0 momentum p ?



Topic: Nuclear and Particle Physics

Sub-Topic: Particle Physics

Ans.: (A)

Solution: The conservation of linear momentum revealed that when decay at rest then angle of produced photon will be 180° . However, when the decayed photon is not at rest then the laws of conservation of momentum again says that that the angle of produced photon will be smaller than 180° . This property is only observable in option (A).

Q.49 A particle has wavefunction

$$\psi(x, y, z) = Nze^{-\alpha(x^2+y^2+z^2)},$$

where N is a normalization constant and α is a positive constant. In this state, which one of the following options represents the eigenvalues of L^2 and L_z respectively?

Some values of Y_l^m are :

$$Y_0^0 = \sqrt{\frac{1}{4\pi}}, Y_1^0 = \sqrt{\frac{3}{4\pi}} \cos \theta, Y_1^{\pm 1} = \mp \sqrt{\frac{3}{8\pi}} \sin \theta e^{\pm i\phi}$$

- (A) 0 and 0 (B) \hbar^2 and $-\hbar$ (C) $2\hbar^2$ and 0 (D) \hbar^2 and \hbar

Topic: Quantum Mechanics

Sub-Topic: Angular momentum

Ans. : (C)

Solution: $\psi(x, y, z) = Nz \exp-(x^2 + y^2 + z^2)$





$$\psi(r, \theta, \phi) = Nr \cos \theta \exp-(r^2) \Rightarrow N \sqrt{\frac{4\pi}{3}} Y_1^0 \exp-(r^2)$$

So $l = 1, m = 0$ measurement of L^2 is $2\hbar^2$ and measurement of L_z is $0\hbar$

Q.50 The wavefunction of a particle in one dimension is given by

$$\psi(x) = \begin{cases} M, & -a < x < a \\ 0, & \text{otherwise} \end{cases}$$

Here M and a are positive constants. If $\varphi(p)$ is the corresponding momentum space wavefunction, which one of the following plots best represents $|\varphi(p)|^2$?



Topic: Quantum Mechanics

Sub-Topic: Wave function

Ans. : (C)

Solution: $\psi(x) = \begin{cases} M = -a < x < a \\ 0, \text{otherwise} \end{cases}$

Normalized wave function is $\psi(x) = \begin{cases} \frac{1}{\sqrt{2a}} = -a < x < a \\ 0, \text{otherwise} \end{cases}$

$$\phi(p) = \frac{1}{\sqrt{2\pi\hbar}} \int_{-\infty}^{\infty} \psi(x) \exp\left(-i \frac{px}{\hbar}\right) dx \Rightarrow \frac{1}{\sqrt{2\pi\hbar}} \int_{-a}^a \frac{1}{\sqrt{2a}} \exp\left(-i \frac{px}{\hbar}\right) dx$$

$$\phi(p) = \sqrt{\frac{\hbar}{\pi a}} \frac{\sin\left(\frac{pa}{\hbar}\right)}{p} \Rightarrow |\phi(p)|^2 = \frac{\hbar}{\pi a} \frac{\sin^2\left(\frac{pa}{\hbar}\right)}{p^2}$$

Q.51 Consider a particle in a two-dimensional infinite square well potential of side L , with $0 \leq x \leq L$ and $0 \leq y \leq L$. The wavefunction of the particle is zero only along the line $y = \frac{L}{2}$, apart from the boundaries of the well. If the energy of the particle in this state is E , what is the energy of the ground state?

- (A) $\frac{1}{4}E$ (B) $\frac{2}{5}E$ (C) $\frac{3}{8}E$ (D) $\frac{1}{2}E$

Topic: Quantum Mechanics

Sub-Topic: Particle in Box

Ans. : (B)

Solution: For two-dimensional box $E_{n_x, n_y} = (n_x^2 + n_y^2) \frac{\pi^2 \hbar^2}{2mL^2}$

For given information $n_x = 1, n_y = 2$, So $E_{1,2} = (1^2 + 2^2) \frac{\pi^2 \hbar^2}{2mL^2} \Rightarrow \frac{\pi^2 \hbar^2}{2mL^2} = \frac{E}{5}$

Ground state is $E_{1,1} = (1^2 + 1^2) \frac{\pi^2 \hbar^2}{2mL^2} = 2 \cdot \frac{E}{5}$

Q.52 Consider two non-identical spin $\frac{1}{2}$ particles labelled 1 and 2 in the spin product state

$\left| \frac{1}{2}, \frac{1}{2} \right\rangle \left| \frac{1}{2}, -\frac{1}{2} \right\rangle$. The Hamiltonian of the system is

$$H = \frac{4\lambda}{\hbar^2} \vec{S}_1 \cdot \vec{S}_2$$

where \vec{S}_1 and \vec{S}_2 are the spin operators of particles 1 and 2, respectively, and λ is a constant with appropriate dimensions. What is the expectation value of H in the above state?

Topic: Atomic, Molecular and Laser Physics

Sub-Topic: Coupling

Ans. : (A)

Solution: The concept of CG Coefficient demonstrated that the above state can be written in the following

$$\text{form } \left| \frac{1}{2}, \frac{1}{2} \right\rangle \left| \frac{1}{2}, -\frac{1}{2} \right\rangle = \frac{1}{\sqrt{2}} |1, 0\rangle + \frac{1}{\sqrt{2}} |0, 0\rangle$$

$$H = \frac{4\lambda}{\hbar^2} S_1 \cdot S_2 = \frac{2\lambda}{\hbar^2} (S^2 - S_1^2 - S_2^2) \Rightarrow E = 2\lambda (S(S+1) - S_1(S_1+1) - S_2(S_2+1))$$

$$S_1 = \frac{1}{2}, S_2 = \frac{1}{2} \Rightarrow S = 1, 0$$

$$\text{Case (1) } S_1 = \frac{1}{2}, S_2 = \frac{1}{2} \& S = 1 \Rightarrow E = \lambda$$

$$\text{Case (2) } S_1 = \frac{1}{2}, S_2 = \frac{1}{2} \& S = 0 \Rightarrow E = -3\lambda$$

$$\langle E \rangle = -\frac{3}{2}\lambda + \frac{1}{2}\lambda = -\lambda$$

Q.53 A spin $\frac{1}{2}$ particle is in a spin up state along the x -axis (with unit vector \hat{x}) and is denoted as $\left| \frac{1}{2}, \frac{1}{2} \right\rangle_x$.

What is the probability of finding the particle to be in a spin up state along the direction \hat{x}' , which lies in the xy -plane and makes an angle θ with respect to the positive x -axis, if such a measurement is made?

- (A) $\frac{1}{2} \cos^2 \frac{\theta}{4}$ (B) $\cos^2 \frac{\theta}{4}$ (C) $\frac{1}{2} \cos^2 \frac{\theta}{2}$ (D) $\cos^2 \frac{\theta}{2}$

Topic: Quantum Mechanics

Sub-Topic: Spin

Ans. : (D)

Solution: $S_x = \frac{\hbar}{2} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ eigen value is $\frac{\hbar}{2}, -\frac{\hbar}{2}$ with corresponding eigen state $\left| \frac{1}{2}, \frac{1}{2} \right\rangle_x = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ 1 \end{pmatrix}$ and

$$\left| \frac{1}{2}, -\frac{1}{2} \right\rangle_x = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ -1 \end{pmatrix}$$

The \vec{S} along \hat{x}' which is making an angle θ with the x axis in the x - y plane is

$$A = \vec{S} \cdot \hat{x}' \text{ is } S_x \cos \theta + S_y \sin \theta \text{ where } \hat{x}' = \cos \theta \hat{i} + \sin \theta \hat{j} \text{ And } S_y = \frac{\hbar}{2} \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}$$

So, operator $A = \begin{bmatrix} 0 & \exp(-i\theta) \\ \exp(i\theta) & 0 \end{bmatrix}$ the eigenvalue of the eigenstate of the operator $\frac{\hbar}{2}, -\frac{\hbar}{2}$

$$\text{with eigenstate } \left| \frac{1}{2}, \frac{1}{2} \right\rangle_{x'} = \frac{1}{\sqrt{2}} \begin{pmatrix} \exp\left(\frac{-i\theta}{2}\right) \\ \exp\left(\frac{i\theta}{2}\right) \end{pmatrix} \text{ and } \left| \frac{1}{2}, -\frac{1}{2} \right\rangle_{x'} = \frac{1}{\sqrt{2}} \begin{pmatrix} -\exp\left(\frac{-i\theta}{2}\right) \\ \exp\left(\frac{i\theta}{2}\right) \end{pmatrix}$$

The probability that state $\left| \frac{1}{2}, \frac{1}{2} \right\rangle_{\hat{x}}$ in eigen state $\left| \frac{1}{2}, \frac{1}{2} \right\rangle_x$ is

$$= \left| \left\langle \frac{1}{2}, \frac{1}{2} \left| \frac{1}{2}, \frac{1}{2} \right\rangle_{\hat{x}} \right. \right|^2 = \left| \frac{1}{\sqrt{2}} (1 \quad 1) \frac{1}{\sqrt{2}} \begin{pmatrix} \exp -\frac{i\theta}{2} \\ \exp \frac{i\theta}{2} \end{pmatrix} \right|^2 = \left| \frac{\exp -\frac{i\theta}{2} + \exp -\frac{i\theta}{2}}{2} \right|^2 = \cos^2 \frac{\theta}{2}$$

The probability that state $\left| \frac{1}{2}, -\frac{1}{2} \right\rangle_{\hat{x}}$ in eigen state $\left| \frac{1}{2}, \frac{1}{2} \right\rangle_x$ is

$$= \left| \left\langle \frac{1}{2}, \frac{1}{2} \left| \frac{1}{2}, -\frac{1}{2} \right\rangle_{\hat{x}} \right. \right|^2 = \left| \frac{1}{\sqrt{2}} (1 \quad 1) \frac{1}{\sqrt{2}} \begin{pmatrix} -\exp -\frac{i\theta}{2} \\ \exp \frac{i\theta}{2} \end{pmatrix} \right|^2 = \left| \frac{-\exp -\frac{i\theta}{2} + \exp -\frac{i\theta}{2}}{2} \right|^2 = \sin^2 \frac{\theta}{2}$$

- Q.54 Different spectral lines of the Balmer series (transitions $n \rightarrow 2$, with n being the principal quantum number) fall one at a time on a Young's double slit apparatus. The separation between the slits is d and the screen is placed at a constant distance from the slits. What factor should d be multiplied by to maintain a constant fringe width for various lines, as n takes different allowed values?

(A) $\frac{n^2 - 4}{4n^2}$ (B) $\frac{n^2 + 4}{4n^2}$ (C) $\frac{4n^2}{n^2 - 4}$ (D) $\frac{4n^2}{n^2 + 4}$

Topic: Optics

Sub-Topic: Interference

Ans. : (C)

Solution: $\lambda = \frac{4n^2}{n^2 - 4}$

Fringe width $\beta = \frac{D\lambda}{d}$ $\frac{4n^2}{n^2 - 4} \Rightarrow d' = k \frac{4n^2}{n^2 - 4}$

- Q.55 Under parity and time reversal transformations, which of the following statements is(are) TRUE about the electric dipole moment \mathbf{p} and the magnetic dipole moment $\boldsymbol{\mu}$?

- (A) \mathbf{p} is odd under parity and $\boldsymbol{\mu}$ is odd under time reversal
- (B) \mathbf{p} is odd under parity and $\boldsymbol{\mu}$ is even under time reversal
- (C) \mathbf{p} is even under parity and $\boldsymbol{\mu}$ is odd under time reversal
- (D) \mathbf{p} is even under parity and $\boldsymbol{\mu}$ is even under time reversal

Topic: Nuclear and Particle Physics

Sub-Topic: Particle Physics

Ans. : (A)

Solution: $p = \sum q_i r_i \Rightarrow \hat{P}: p = \sum q_i (-r_i) = -p$ (odd)

$$\mu = -\frac{e}{2m} L \Rightarrow \hat{T} : \mu = -\frac{e}{2m} (-L) = -\mu \text{ (odd)}$$

Q.56 Consider the complex function

$$f(z) = \frac{z^2 \sin z}{(z - \pi)^4}$$

At $Z = \pi$, which of the following options is(are) CORRECT?

- (A) The order of the pole is 4 (B) The order of the pole is 3
 (C) The residue at the pole is $\frac{\pi}{6}$ (D) The residue at the pole is $\frac{2\pi}{3}$

Topic: Mathematical Physics

Sub-Topic:

Complex

Analysis

Ans. : (B)

Solution: Put $z - \pi = t \Rightarrow f(z) = \frac{(t + \pi)^2 \sin(t + \pi)}{t^4} = \frac{(t + \pi)^2 (-\sin t)}{t^4}$

$$= \frac{-(t^2 + \pi^2 + 2t\pi) \left(t - \frac{t^3}{3!} + \dots \right)}{t^4}$$

So, Pole is of order 3. Residue = Coefficient of $\frac{1}{t} = \frac{1}{6}(\pi^2 - 6)$

Q.57 Consider the vector field \vec{V} consisting of the velocities of points on a thin horizontal disc of radius $R = 2$ m, moving anticlockwise with uniform angular speed $\omega = 2$ rad/sec about an axis passing through its center. If $V = |\vec{V}|$, then which of the following options is(are) CORRECT? (In the options, \hat{r} and $\hat{\theta}$ are unit vectors corresponding to the plane polar coordinates r and θ).

You may use the fact that in cylindrical coordinates (S, ϕ, z) (S is the distance from the Z -axis), the gradient, divergence, curl and Laplacian operators are:

$$\vec{\nabla} f = \frac{\partial f}{\partial s} \hat{s} + \frac{1}{s} \frac{\partial f}{\partial \phi} \hat{\phi} + \frac{\partial f}{\partial z} \hat{z};$$

$$\vec{\nabla} \cdot \vec{A} = \frac{1}{s} \frac{\partial}{\partial s} (sA_s) + \frac{1}{s} \frac{\partial A_\phi}{\partial \phi} + \frac{\partial A_z}{\partial z}$$

$$\vec{\nabla} \times \vec{A} = \left(\frac{1}{s} \frac{\partial A_z}{\partial \phi} - \frac{\partial A_\phi}{\partial z} \right) \hat{s} + \left(\frac{\partial A_s}{\partial z} - \frac{\partial A_z}{\partial s} \right) \hat{\phi} + \frac{1}{s} \left(\frac{\partial}{\partial s} (sA_\phi) - \frac{\partial A_s}{\partial \phi} \right) \hat{z}$$

$$\vec{\nabla}^2 f = \frac{1}{s} \frac{\partial}{\partial s} \left(s \frac{\partial f}{\partial s} \right) + \frac{1}{s^2} \frac{\partial^2 f}{\partial \phi^2} + \frac{\partial^2 f}{\partial z^2}$$

- (A) $\vec{\nabla} V = 2\hat{r}$
 (B) $\vec{\nabla} \cdot \vec{V} = 2$
 (C) $\vec{\nabla} \times \vec{V} = 4\hat{z}$, where \hat{z} is a unit vector perpendicular to the (r, θ) plane

(D) $\vec{\nabla}^2 V = \frac{4}{3}$ at $r = 1.5$ m

Topic: EMT

Sub-Topic: Magnetostatic

Ans. : (A), (C) and (D)

Solution: $\vec{v} = \vec{\omega} \times \vec{r}$, $\vec{V} = \omega \hat{z} \times r \hat{r} = \omega r \hat{\phi}$, $|\vec{v}| = \omega r$

(A) $\vec{\nabla} v = \frac{\partial}{\partial r} = \frac{\partial}{\partial r} (\omega r) = \omega \hat{r} = 2\hat{r} \therefore \omega = 2 \text{ rad/se}$

(B) $\vec{\nabla} \cdot \vec{v} = \frac{1}{r} \frac{\partial}{\partial \phi} (r\omega) = 0$ (solenoidal)

(C) $\nabla \times v = \frac{1}{r} \begin{vmatrix} \hat{r} & r\hat{\phi} & z \\ \frac{\partial}{\partial r} & \frac{\partial}{\partial \phi} & \frac{\partial}{\partial z} \\ 0 & v & 0 \end{vmatrix}$, $\hat{v} = r\omega \hat{\phi}$

$= \frac{\hat{r}}{r} \left[\frac{\partial}{\partial \phi} (0) - \frac{\partial}{\partial r} (v) \right] - r\hat{\phi} \left[\frac{\partial}{\partial r} (0) - \frac{\partial}{\partial r} (0) \right] + \hat{z} \left[\frac{\partial}{\partial r} (rv) - \frac{\partial}{\partial \phi} (0) \right] = 4\hat{z}$

(D) $\nabla^2 v = \frac{1}{r} \frac{\partial}{\partial r} \left(r \frac{\partial f}{\partial r} \right) + \frac{1}{r^2} \frac{\partial^2 f}{\partial \phi^2} + \frac{\partial^2 f}{\partial z^2} = \frac{1}{r} \frac{\partial}{\partial r} (r\omega) = \frac{\omega}{r} = \frac{2 \times 2}{3} = \frac{4}{3}$, at $r = 1.5$

Q.58 A slow moving π^- particle is captured by a deuteron (d) and this reaction produces two neutrons (n) in the final state, i.e., $\pi^- + d \rightarrow n + n$. Neutron and deuteron have even intrinsic parities, whereas π^- has odd intrinsic parity. L and S are the orbital and spin angular momenta, respectively of the system of two neutrons. Which of the following statements regarding the final two-neutron state is(are) CORRECT?

- (A) It has odd parity
- (B) $L + S$ is odd
- (C) $L = 1, S = 1$
- (D) $L = 2, S = 0$

Topic: Nuclear and Particle Physics

Sub-Topic: Particle Physics

Ans.: (A) & (C)

Solution: $\pi^- + d \rightarrow n + n$

The slow-moving means $L = 0$ for π^- . The deuteron is having 3_{S_1} & 3_{D_1} which represents $L = \text{Even}$ for it. Also, it is important to be mentioned here that deuteron is fermion having intrinsic parity +1.

From the conservation of Parity in strong interaction

$(-1)^{L'} \pi_n \pi_d = \pi_n \pi_n (-1)^L$

$(-)(+) = (+)(+)(-1)^L$ (Since, $L' = \text{Even}$)

$$(-)^L = (-1)^L$$

Thus, $L = \text{Odd} = 1, 3, 5, \dots$

The conservation of parity itself says the product is having odd parity. Thus, option (A) is correct.

From spin conservation


$$\pi^- + d \rightarrow n + n$$

$$S \quad 0 + 1 \quad \frac{1}{2} + \frac{1}{2}$$

Resultant S of product will be 1

So, spin of final two neutron state option (C) is correct and option (D) is wrong. $L + S$ will be even.

- Q.59 Two independent electrostatic configurations are shown in the figure. Configuration (I) consists of an isolated point charge $q = 1\text{C}$, and configuration (II) consists of another identical charge surrounded by a thick conducting shell of inner radius $R_1 = 1\text{ m}$ and outer radius $R_2 = 2\text{ m}$, with the charge being at the center of the shell. $W_I = \frac{\epsilon_0}{2} \int E_I^2 dV$ and $W_{II} = \frac{\epsilon_0}{2} \int E_{II}^2 dV$, where E_I and E_{II} are the magnitudes of the electric fields for configurations (I) and (II) respectively, ϵ_0 is the permittivity of vacuum, and the volume integrations are carried out over all space. If $8\pi\epsilon_0|W_I - W_{II}| = \frac{1}{n}$, what is the value of the integer n ?



Topic: EMT

Sub-Topic: Electrostatic Energy

Ans.:

Solution:

$$W = \frac{\epsilon_0}{2} \int E^2 d\tau, q = 1\text{C}$$

$$R_1 = 1\text{ m}, R_2 = 2\text{ m}$$

$$\frac{8\pi}{\epsilon_0} [W_I - W_{II}] = \frac{1}{n}$$

$$W_I = \frac{\epsilon_0}{2} \int_{\epsilon}^{\infty} \left(\frac{1}{4\pi\epsilon_0} \frac{q}{r^2} \right)^2 4\pi r^2 dr = \frac{1}{8\pi\epsilon_0} \frac{1}{\epsilon}$$

$$W_{II} = \frac{\epsilon_0}{2} \left[\int_{\epsilon}^{R_1} \left(\frac{1}{4\pi\epsilon_0} \frac{q}{r^2} \right)^2 4\pi r^2 dr + 0 + \int_{R_2}^{\infty} \left(\frac{1}{4\pi\epsilon_0} \frac{q}{r^2} \right)^2 4\pi r^2 dr \right]$$

$$W_I - W_{II} = \frac{1}{8\pi\epsilon} \times \frac{1}{2}, \quad 8\pi\epsilon(W_I - W_{II}) = \frac{1}{2}, \quad n=2$$

Q.60 In pion nucleon scattering, the pion and nucleon can combine to form a short-lived bound state called the Δ particle ($\pi + N \rightarrow \Delta$). The masses of the pion, nucleon and the Δ particle are $140\text{MeV}/c^2$, $938\text{MeV}/c^2$ and $1230\text{MeV}/c^2$, respectively. In the lab frame, where the nucleon is at rest, what is the minimum energy (in MeV/c^2 , rounded off to one decimal place) of the pion to produce the Δ particle?

Topic: Nuclear and Particle Physics

Sub-Topic: Nuclear Reaction

Ans. : 340 MeV (Not matched)

$$\text{Solution: } E_{\pi} = \frac{m_{\Delta}^2 - m_N^2 - m_{\pi}^2}{2m_N} = \frac{1240^2 - 938^2 - 140^2}{2 \times 938} = 340\text{ MeV}$$

Q.61 Consider an electromagnetic wave propagating in the z-direction in vacuum, with the magnetic field given by $\vec{B} = \vec{B}_0 e^{i(kz - \omega t)}$. If $B_0 = 10^{-8}\text{ T}$, the average power passing through a circle of radius 1.0 m placed in the xy plane is P (in Watts). Using $\epsilon_0 = 10^{-11} \frac{\text{C}^2}{\text{Nm}^2}$, what is the value of $\frac{10^3 P}{\pi}$ (rounded off to one decimal place)?

Topic: EMT

Sub-Topic: Propagation of EM Wave

Ans. : 13.5

$$\begin{aligned} \text{Solution: Intensity } I &= \frac{P}{A} = \langle u \rangle \cdot c = \frac{B_0^2}{2\mu_0} C = \frac{B_0^2 C}{2} \times \epsilon_0 C^2 \\ \Rightarrow P &= \frac{B_0^2 C^3 \epsilon_0}{2} \times \pi(1)^2 \Rightarrow \frac{10^3 P}{\pi} = \frac{10^3 B_0^2 C^3 \epsilon_0}{2} \\ \Rightarrow \frac{10^3 P}{\pi} &= \frac{10^3}{2} \times (10^{-8})^2 (3 \times 10^8)^3 \times 10^{-1} \\ &= \frac{10^3}{2} \times 10^{-16} \times 27 \times 10^{24} \times 10^{-1} = \frac{27}{2} = 13.5 \end{aligned}$$

Q.62 An α -particle is emitted from the decay of Americium (Am) at rest, i.e., ${}_{94}^{241}\text{Am} \rightarrow {}_{92}^{237}\text{U} + \alpha$. The rest masses of ${}_{94}^{241}\text{Am}$, ${}_{92}^{237}\text{U}$ and α are $224.544 \text{ GeV}/c^2$, $220.811 \text{ GeV}/c^2$ and $3.728 \text{ GeV}/c^2$ respectively. What is the kinetic energy (in MeV/c^2 , rounded off to two decimal places) of the α -particle?

Topic: Nuclear and Particle Physics

Sub-Topic: Radioactivity

Ans. : 4.917

Solution: ${}_{94}^{241}\text{Am} \rightarrow {}_{92}^{237}\text{U} + \alpha \Rightarrow Q_\alpha = [M_{Am} - M_U - M_\alpha]c^2$
 $= [224.544 - 220.811 - 3.728] \frac{\text{GeV}}{c^2} \times c^2 = 0.005 \text{ GeV} = 5 \text{ MeV}$

$$K_\alpha = \frac{A-4}{4} Q_\alpha = \frac{241-4}{241} \times 5 \text{ MeV} = 4.917 \text{ MeV}$$

Q.63 Consider 6 identical, non-interacting, spin $\frac{1}{2}$ atoms arranged on a crystal lattice at absolute temperature T . The Z -component of the magnetic moment of each of these atoms can be $\pm\mu_B$. If P and Q are the probabilities of the net magnetic moment of the solid being $2\mu_B$ and $6\mu_B$ respectively, what is the value of $\frac{P}{Q}$ (in integer)

Topic: Thermodynamic and Statistical Mechanics

Sub-Topic: Statistical Distribution

Ans. : 63: 15

Solution: For total moment to be $2\mu_B$ the arrangement is like $\uparrow\uparrow\uparrow\uparrow\downarrow\downarrow \rightarrow 2\mu_B \Rightarrow P = \frac{{}^6C_4}{2^6}$

For total moment to be $6\mu_B$ the arrangement is like $\uparrow\uparrow\uparrow\uparrow\uparrow\uparrow \Rightarrow Q = \frac{{}^6C_6}{2^6}$

$$\frac{P}{Q} = \frac{6!}{2! \cdot 4!} = \frac{6 \times 5}{2} = 15$$

Q.64 Two identical, non-interacting ${}^4\text{He}_2$ atoms are distributed among 4 different nondegenerate energy levels. The probability that they occupy different energy levels is p . Similarly, two ${}^3\text{He}_2$ atoms are distributed among 4 different non-degenerate energy levels, and the probability that they occupy different levels is q . What is the value of $\frac{p}{q}$ (rounded off to one decimal place)?

Topic: Thermodynamic and Statistical Mechanics

Sub-Topic: Statistical distribution

Ans. : 64: 0.6

Solution: Bosons, $p = 6/10$

| | | | | | | | | | |
|----|----|----|----|---|---|---|---|---|---|
| AA | | | | A | | | A | | A |
| | | | | | | | | | |
| | AA | | | A | A | | | A | |
| | | AA | | | A | A | A | | |
| | | | AA | | | A | | A | A |

Fermions, $p = 6/6$

| | | | | | |
|---|---|---|---|---|---|
| A | | | A | | A |
| A | A | | | A | |
| | A | A | A | | |
| | | A | | A | A |

$p/q = 0.6$

Q.65 Two identical bodies kept at temperatures 800 K and 200 K act as the hot and the cold reservoirs of an ideal heat engine, respectively. Assume that their heat capacity (C) in Joules/ K is independent of temperature and that they do not undergo any phase change. Then, the maximum work that can be obtained from the heat engine is $n \times C$ Joules. What is the value of n (in integer)?

Topic: Thermodynamic and Statistical Mechanics

Sub-Topic: Thermodynamic Process

Ans. : 200

Solution: $Q_1 = C[T_1 - T']$ & $Q_2 = C[T' - T_2]$

$$W = Q_1 - Q_2 = C[T_1 - T' - T' + T_2] = C[T_1 + T_2 - 2T']$$

$$\Delta S = \Delta S_1 + \Delta S_2 = 0 \Rightarrow \int_{T_1}^{T'} \frac{CdT}{T} + \int_{T_2}^{T'} \frac{CdT}{T} = 0 \Rightarrow C \ln \frac{T'}{T_1} + C \ln \frac{T'}{T_2} = 0$$

$$\Rightarrow T' = \sqrt{T_1 T_2} \Rightarrow W = [800 + 200 - 2\sqrt{1600}] = 200$$