

JNU MSc 2019

Q1. Longitudinal waves are

- (a) Plane polarized (b) Circularly polarized
(c) Elliptically polarized (d) Unpolarized

Ans. : (d)

Q2. One nanometer is equal to

- (a) 0.1 \AA (b) 10 \AA (c) 100 \AA (d) 1000 \AA

Ans. : (b)

Q3. According to the Dulong-Petit law, the atomic heat which is a product of atomic weight and specific heat, of most of the elements in solid state

- (a) Is constant (b) increases with atomic number
(c) Decrease with atomic number (d) Does not depend on atomic weight

Ans. : (a)

Q4. An X-ray beam consists of

- (a) Electrons (b) Protons (c) Neutrons (d) Photons

Ans. : (d)

Q5. A thermocouple is a device to measure

- (a) Pressure (b) Volume (c) Density (d) Temperature

Ans. : (d)

Q6. What would be the frequency of the photon produced when an electron of energy 20 keV is brought to rest in a collision with a heavy nucleus?

- (a) $4.84 \times 10^{18} \text{ Hz}$ (b) $5 \times 10^{18} \text{ Hz}$
(c) $4.23 \times 10^{18} \text{ Hz}$ (d) $3.84 \times 10^{18} \text{ Hz}$

Ans. : (a)

Q7. Consider a planet of mass m , in circular motion with angular momentum, L . The planet orbits a star of mass, M and the orbit radius is r . If the radius of the orbit is changed from r to $\frac{r}{2}$, what would be the new value of angular momentum?

- (a) L (b) $L/2$ (c) $\frac{L}{\sqrt{2}}$ (d) $\sqrt{2}L$

Ans. : (c)

Q8. At time $t=0$, a series RC circuit is connected to an emf of $9V$. How long will it take for the capacitor to reach $8V$?

- (a) RC (b) $\frac{1}{RC}$ (c) $RC \ln 9$ (d) $\ln 9$

Ans. : (c)

Q9. Which of the following quantities has the same physical dimension as that of $\frac{h}{e^2}$, where h is Planck's constant and e is electronic charge?

- (a) Magnetic flux (b) Electrical resistance
(c) Magnetic field (d) Electrical resistivity

Ans. : (b)

Q10. For a hydrogen atom the spacing between successive energy levels is given by $\Delta_n = E_{n+1} - E_n$, where n is the quantum number. Which of the following statements is true?

- (a) Δ_n is constant
(b) Δ_n increases as n increases
(c) Δ_n decreases as n increases
(d) Δ_n increases and then decreases with n

Ans. : (c)

Q11. Consider a momentum conservation experiment where two masses m_1 and m_2 are collided head-on with velocities v_1 and v_2 , respectively, the measured values are $m_1 = 200 \pm 2g$, $v_1 = 5.5 \pm 0.1 m/s$ and $v_2 = 10 \pm 0.4 m/s$. What is the fractional error associated with mass m_2 of the other body

- (a) ± 7.7 (b) ± 0.77 (c) ± 10.1 (d) ± 0.07

Ans. : (a)

Q12. A sinusoidal wave moving along a string in the x -direction is described by

$$y(x,t) = 0.002 \sin(10x - 120t)$$

What is the propagation speed of the wave?

- (a) $12 m/s$ (b) $10 m/s$ (c) $120 m/s$ (d) $1200 m/s$

Ans. : (a)

Q13. The black body radiation emitted from a cavity of volume V at temperature T has chemical potential equal to (N is the number of photons emitted)

- (a) N (b) 0 (c) $\frac{1}{T}$ (d) $\frac{V}{T}$

Ans. : (b)

Q14. A $100W$ electric bulb has an efficiency of 2.5% . Assuming it is a point source, the intensity at a distance of $3m$ will be

- (a) $2.5W/m^2$ (b) $25W/m^2$
(c) $0.025W/m^2$ (d) $0.022W/m^2$

Ans. : (d)

Q15. An electron has a speed of $300m/s$, accurate to 0.01% . With what accuracy can we determine the position of the electron? (mass of electron = $9.1 \times 10^{-31} kg$, Planck's constant = $6.6 \times 10^{-34} J \cdot s$)

- (a) $2.4 nm$ (b) $2.4 \mu m$ (c) $2.4 mm$ (d) $2.4 cm$

Ans. : (d)

Q16. A burst of 10^{14} electrons uniformly accelerated to an energy of $15 MeV$ is stopped by a copper target block of mass $100 g$. Assuming the block is thermally insulated, what is the rise in its temperature? (specific heat of copper is $0.09 cal/g K$)

- (a) $6.3 K$ (b) $0.4 K$ (c) $1.7 K$ (d) $5.1 K$

Ans. : (a)

Q17. The function $y = ax^2 - bx + c$, where a, b and c are positive and constants, has a minima at $x =$

- (a) $\frac{b}{2a}$ (b) $\frac{a}{2b}$ (c) $\frac{b}{a}$ (d) $\frac{a}{b}$

Ans. : (a)

Q18. During radioactive decay a nucleus emits a gamma ray with energy of $1.35 MeV$. What is the wavelength of this photon?

- (a) $920 fm$ (b) $920 nm$ (c) $920 pm$ (d) 920 \AA

Ans. : (a)

Q19. The adiabatic compressibility of an ideal gas is equal to (P is pressure and V is volume)

- (a) $\frac{1}{P}$ (b) $\frac{P}{V}$ (c) P (d) $\frac{V}{P}$

Ans. : (a)

Q20. The angle between the vectors $\vec{a} = \hat{i} + \hat{j}$ and $\vec{b} = \hat{i} + \hat{j} + \hat{k}$ is

- (a) 0° (b) 45° (c) $\cos^{-1}\left(\frac{1}{3}\right)$ (d) $\cos^{-1}\left(\sqrt{\frac{2}{3}}\right)$

Ans. : (d)

Q21. A 2 mW laser light is emitted at a frequency of $6 \times 10^{14}\text{ Hz}$. How many photons on average are emitted by this source per second? (Plank's constant $6.6 \times 10^{-34}\text{ J}\cdot\text{s}$)

- (a) 1×10^{15} (b) 2×10^{15} (c) 3×10^{15} (d) 5×10^{15}

Ans. : (d)

Q22. A particle of mass m moves in a circle of radius r with uniform angular speed ω . The work done by the centripetal force in half of a complete rotation is

- (a) 0 (b) $2\pi m\omega^2 r^2$ (c) $\frac{\pi m\omega^2 r^2}{2}$ (d) $2\pi m\omega^2$

Ans. : (a)

Q23. Resistances R_1 and R_2 are connected in parallel and I is the total current flowing in the circuit. I_1 is the current flowing through R_1 . Which of the following conditions will produce minimum joule heating in the circuit?

- (a) $I_1 = I \left(\frac{R_2}{R_1 + R_2} \right)$ (b) $I_1 = I_2 \left(\frac{R_2}{R_1 + R_2} \right)$
 (c) $I_1 = I_2 \left(\frac{R_2}{R_1} \right)$ (d) $I_1 = I_2 \left(\frac{R_1}{R_2} \right)$

Ans. : (a)

Q24. In a two-level atom, the energy gap is E . The probability of finding the atom in the excited state at temperature T will be

- (a) $\exp\left(-\frac{E}{k_B T}\right)$ (b) $\frac{1}{1 + \exp\left(-\frac{E}{k_B T}\right)}$
 (c) $\frac{\exp\left(\frac{E}{k_B T}\right)}{1 + \exp\left(-\frac{E}{k_B T}\right)}$ (d) $\frac{\exp\left(-\frac{E}{k_B T}\right)}{1 + \exp\left(-\frac{E}{k_B T}\right)}$

Ans. : (d)

Q25. Consider a two-dimensional quantum harmonic oscillator with frequency ω . How many energy levels are there with energy $11\hbar\omega$?

- (a) 5 (b) 8 (c) 11 (d) 21

Ans. : (c)

Q26. What is the entropy change when 1kg of ice at $0^{\circ}C$ melts reversibly to water at the same temperature? (latent heat of melting of ice = 79.6 cal/g)

- (a) $122\text{ kJ}\cdot\text{K}^{-1}$ (b) $12.2\text{ kJ}\cdot\text{K}^{-1}$ (c) $1.22\text{ kJ}\cdot\text{K}^{-1}$ (d) $0.122\text{ kJ}\cdot\text{K}^{-1}$

Ans. : (c)

Q27. The equation of motion of a particle of mass m in one-dimension is

$$m \frac{d^2x}{dt^2} = -ax - 3bx^2 - 4cx^3$$

where a, b and c are constants of appropriate dimension. The quantity that remains constant during its motion is

- (a) $\frac{1}{2}m\dot{x}^2 + \frac{1}{2}ax^2 + bx^3 + cx^4$ (b) $\frac{1}{2}m\dot{x}^2 + ax^2 + bx^3 + cx^4$
 (c) $\frac{1}{2}m\dot{x}^2 + \frac{1}{2}ax^2 + \frac{1}{3}bx^3 + cx^4$ (d) $\frac{1}{2}m\dot{x}^2 + ax^2 + \frac{1}{3}bx^3 + \frac{1}{4}cx^4$


Ans. : (a)

Q28. The crystal structure of CsCl is a simple cubic lattice. Each unit cell of CsCl will contain

- (a) 1 atom (b) 2 atoms (c) 3 atoms (d) 4 atoms

Ans. : (b)

Q29. The reading in the ammeter A is



- (a) 0.5454 A (b) 5.5450 A (c) 5.4555 A (d) 1.5455 A

Ans. : (a)

Q30. An ideal gas undergoes isothermal expansion at temperature T from volume V_1 to V_2 . The entropy change per mole is

- (a) $R \left(\frac{V_2}{V_1} \right)$ (b) $R \left(\frac{V_1}{V_2} \right)$ (c) $R \ln \left(\frac{V_2}{V_1} \right)$ (d) $R \ln \left(\frac{V_1}{V_2} \right)$

Ans. : (c)

Q31. Which of the following is responsible for the existence of the Fermi surface in metals?

- (a) Nuclear force
 (b) Coulomb repulsion between electrons
 (c) Bose-Einstein condensation
 (d) Pauli exclusion principle

Ans. : (d)

Q32. A sodium vapour lamp emits yellow light corresponding to two wavelengths 589 and 589.59 nm. What is the minimum number of rulings must a diffraction grating have to resolve these two lines in the first order?

- (a) 589 (b) 700 (c) 900 (d) 1000

Ans. : (d)

Q33. If $z = x + iy$, the value of $|\sin z|^2$ is

- (a) $\sin^2 x + \sin^2 y$ (b) $\sin^2 x + \cos^2 y$
 (c) $\sin^2 x + \sinh^2 y$ (d) $\sin^2 x + \cosh^2 y$

Ans. : (a)

Q34. If a signal passing through a gate is inhibited by sending a LOW into one of the inputs, and the output is HIGH, the gate is

- (a) an AND gate (b) a NAND gate
 (c) a NOR gate (d) an OR gate

Ans. : (b)

Q35. If \hbar is the reduced Planck's constant, c is the speed of light, and G is the universal gravitational constant, which of the following has the dimension of length?

- (a) $\frac{\hbar G}{c^2}$ (b) $\sqrt{\frac{\hbar c}{8\pi G}}$ (c) $\sqrt{\frac{\hbar G}{c^5}}$ (d) $\sqrt{\frac{\hbar G}{c^3}}$

Ans. : (d)